


Journal Article

**Brain lymphoma: usefulness of the magnetic resonance spectroscopy**

Journal	Journal of Neuro-Oncology
Publisher	Springer Netherlands
ISSN	0167-594X (Print) 1573-7373 (Online)
Issue	Volume 86, Number 2 / January, 2008
Category	Clinical-Patient Studies
DOI	10.1007/s11060-007-9468-2
Pages	225-229
Subject Collection	Medicine
SpringerLink Date	Wednesday, September 05, 2007

Sophie Taillibert¹, Rémy Guillevin^{2, 3} , Carole Menuel², Marc Sanson¹, Khê Hoang-Xuan¹, Jacques Chiras² and Hugues Duffau⁴

(1) Department of Neuro-oncology, Pitié Salpêtrière Hospital, Paris, France

(2) Department of Neuroradiology, Pitié Salpêtrière Hospital, 47-83 boulevard de l'hôpital, Paris, 75651, France

(3) Laboratoire d'imagerie fonctionnelle, INSERM U678, Paris-6 University, Paris, France

(4) Department of Neurosurgery, Gui de Chauliac Hospital, Montpellier, France

Received: 14 June 2007 **Accepted:** 13 August 2007 **Published online:** 4 September 2007

Abstract The diagnosis of primary central nervous system lymphoma (PCNSL) should always be considered as an emergency because of the therapeutic consequences it implies. In immunocompetent patients, it relies on stereotactic biopsy. Unfortunately, clinical and radiological features may be misleading and delay the diagnostic procedure. The case we report here illustrates the contribution of magnetic resonance spectroscopy in the diagnostic approach of a very atypical PCNSL.

Keywords Demyelination - Multiple sclerosis - Magnetic resonance spectroscopy - Primary CNS lymphoma

 **Rémy Guillevin**
Email: remy.guillevin@psl.aphp.fr

References secured to subscribers.