REVIEW



The role of [¹⁸F]fluorodopa positron emission tomography in grading of gliomas

Joy R. Roach^{1,2} · Puneet Plaha^{2,3} · Daniel R. McGowan^{1,4} · Geoff S. Higgins^{1,5}

Received: 13 September 2022 / Accepted: 19 October 2022 $\ensuremath{\textcircled{}}$ The Author(s) 2022

Abstract

Purpose Gliomas are the most commonly occurring brain tumour in adults and there remains no cure for these tumours with treatment strategies being based on tumour grade. All treatment options aim to prolong survival, maintain quality of life and slow the inevitable progression from low-grade to high-grade. Despite imaging advancements, the only reliable method to grade a glioma is to perform a biopsy, and even this is fraught with errors associated with under grading. Positron emission tomography (PET) imaging with amino acid tracers such as [¹⁸F]fluorodopa (¹⁸F-FDOPA), [¹¹C]methionine (¹¹C-MET), [¹⁸F] fluoroethyltyrosine (¹⁸F-FET), and ¹⁸F-FDOPA are being increasingly used in the diagnosis and management of gliomas. **Methods** In this review we discuss the literature available on the ability of ¹⁸F-FDOPA-PET to distinguish low- from high-grade in newly diagnosed gliomas.

Results In 2016 the Response Assessment in Neuro-Oncology (RANO) and European Association for Neuro-Oncology (EANO) published recommendations on the clinical use of PET imaging in gliomas. However, since these recommendations there have been a number of studies performed looking at whether ¹⁸F-FDOPA-PET can identify areas of high-grade transformation before the typical radiological features of transformation such as contrast enhancement are visible on standard magnetic resonance imaging (MRI).

Conclusion Larger studies are needed to validate ¹⁸F-FDOPA-PET as a non-invasive marker of glioma grade and prediction of tumour molecular characteristics which could guide decisions surrounding surgical resection.

Keywords Gliomas · Glioma grading · FDOPA PET · Fluorodopa pet · Brain tumours

	Joy R. Roach Joy.roach@oncology.ox.ac.uk
	Puneet Plaha Puneet.plaha@ouh.nhs.uk
	Daniel R. McGowan Daniel.mcgowan@oncology.ox.ac.uk
	Geoff S. Higgins Geoffrey.higgins@oncology.ox.ac.uk
1	Department of Oncology, University of Oxford, Oxford OX3 7DQ, UK
2	Department of Neurosurgery, Oxford University Hospital NHS FT, John Radcliffe Hospital, L3 West Wing, Oxford OX3 9DU, UK
3	Nuffield Department of Surgical Sciences, University of Oxford, Oxford OX3 7DQ, UK
4	Department of Medical Physics and Clinical Engineering, Oxford University Hospital NHS FT, Churchill Hospital, Oxford OX3 7LE, UK
5	Department of Oncology, Oxford University Hospitals NHS FT, Oxford, UK

Introduction

Glioma is the most common primary brain tumour occurring in adults for which there is no cure. Gliomas are traditionally dichotomised by their grade into low-grade gliomas (LGG) which include grade I and II and high-grade gliomas (HGG) which includes grade III and IV tumours. The most recent publication of the fifth edition of the WHO classification of brain tumours incorporates new information on histological and molecular features into a layered integrated diagnosis and also introduces grading on a 'within tumour type' method [1].

The majority of patients with a low-grade glioma will present acutely with a seizure and as a result have a plain or contrast-enhanced computed tomography (CT) brain scan which identifies an area of abnormality that then requires further delineation with magnetic resonance imaging (MRI). The minimum MRI sequences performed as part of the diagnostic workup for a suspected brain tumour include T2 weighted, fluid attenuated inversion recovery (FLAIR), T1 pre- and post-contrast and diffusion weighted sequences (DWI). The characteristic appearances of low-grade gliomas on MRI sequences will grossly depend on the grade and histological type of glioma. A provisional diagnosis can be made on imaging alone i.e. distinguishing between glioma versus metastasis, and within gliomas in distinguishing lowgrade versus high-grade depending on a number of tumour characteristics [2]. However, there is diagnostic uncertainty about the prediction of WHO grade, and imaging alone is often inaccurate.

The distinction between low- and high-grade glioma on MRI is based on contrast enhancement from a disrupted blood brain barrier and to some extent from diffusion characteristics but this is not always truly predictive of grade and there is often diagnostic uncertainty. The confirmation of grade is normally made on biopsy sampling. However, sampling errors are not uncommon and up to one third of high-grade gliomas may not display the typical imaging characteristics of a high-grade glioma with enhancement [3]. Under grading is particularly associated with large heterogenous tumours and has been reported in 28% – 68% of cases [4–6]. With the introduction of molecular characterisation the risk of sample bias has reduced dramatically as the molecular markers are volume independent but not completely eliminated [7].

Additional imaging techniques and sequences are frequently used to provide additional information to help distinguish the type, grade of tumour and predict transformation. These include positron emission tomography (PET), single photon emission computed tomography (SPECT), MR perfusion and MR spectroscopy (MRS). Gliomas are notoriously heterogeneous in nature and as a result the use of MRS will demonstrate spectra that vary significantly depending on the region sampled [8]. The use of MR perfusion to detect grade has been demonstrated with varying results [9–11]. SPECT imaging has the disadvantage over PET of a lower resolution [12]. PET using amino acid tracers is becoming increasingly common to differentiate gliomas from metastases or other types of tumours [13]. The most commonly used amino acid PET tracers described are [¹¹C] methionine (¹¹C-MET), [¹⁸F]fluoroethyltyrosine (¹⁸F-FET) and [¹⁸F]fluorodopa (¹⁸F-FDOPA). ¹¹C-MET has a short half-life of 20 min which limits its use to centres that have an onsite cyclotron. In comparison ¹⁸F-FET and ¹⁸F-FDOPA have much longer half-lives of 110 min making these tracers more available for clinical use. ¹⁸F-FDOPA demonstrates greater contrast for lesions outside of the striatum when compared to¹⁸F-FET [14]. A type of novel MRI technique, called oxygen enhanced MRI (OEMRI), has the ability to detect areas of hypoxia within solid tumours which if used within glioma could potentially detect areas of high-grade tumour and therefore affect prognosis if detected earlier [15].

The distinction between high-grade and low-grade glioma is important as both entities confer very different prognoses and management strategies. All treatment options aim to prolong survival, maintain quality of life and slow the inevitable progression from low- to high-grade. For both low- and high-grade gliomas the NICE guidelines recommend consideration of maximum safe gross total resection to confirm the histological and molecular diagnosis and for tumours that are not surgically resectable to perform a biopsy [16]. Following surgery, the standard treatment regime for grade IV gliomas (glioblastoma) is radiotherapy with concomitant temozolomide followed by adjuvant temozolomide. In patients over the age of 70 treatment for grade IV gliomas is hypofractionated chemoradiotherapy [16]. Even with full treatment the median survival for grade IV glioblastomas is 12-24 months [17]. For grade II gliomas following surgery, oncological therapy is dependent on the patient's age, extent of resection, 1p/19q codeletion presence and isocitrate dehydrogenase (IDH) mutation status and consists of a combination of radiotherapy and PCV (procarbazine, CCNU [lomustine] and vincristine) chemotherapy [16] (Fig. 1). For grade III gliomas with a 1p/19q codeletion (anaplastic oligodendroglioma) treatment is similar with radiotherapy and PCV chemotherapy. Grade III tumours without a 1p/19q codeletion (anaplastic astrocytoma) require radiotherapy followed by adjuvant temozolomide. The use of non-invasive imaging parameters to accurately detect glioma grade could aid preoperative clinical decision-making when considering biopsy versus resection, extent of resection and timing of surgery.

Current recommendations

In 2016, the European Association of Nuclear Medicine (EANM), European Association of Neurooncology (EANO) and the Response Assessment in Neurooncology (RANO) working group jointly published guidelines on the role of amino acid PET for imaging in gliomas [18, 19]. These evidence-based guidelines recommended a clinical role for the tracer ¹⁸F-FDOPA in the differentiation of glioma recurrence from treatment-induced changes, assessment of treatment response and assessment of prognosis. The recommendations reviewed the literature on the amino acid tracers ¹¹C-MET, ¹⁸F-FET and ¹⁸F-FDOPA in glioma. Multiple studies have found that response to treatment is indicated on PET imaging by a decrease in amino acid tracer uptake with or without a reduction in the volume of metabolically active tumour [20–22] and ¹⁸F-FDOPA has been shown to demonstrate response to bevacizumab therapy better than conventional MRI [23, 24]. Treatment of gliomas with radiotherapy and/or chemotherapy can result in treatment-related changes. There is a temporary alteration in the blood brain



Fig. 1 NICE guidelines for management of glioma following biopsy or surgical resection

barrier (BBB) resulting in contrast enhancement on MRI imaging which mimics tumour progression and is called pseudoprogression. Pseudoprogression typically occurs within 12 weeks of completion of treatment [25, 26]. Differentiating pseudoprogression and radionecrosis from true tumour progression can be challenging with conventional MRI alone and often additional imaging techniques such as amino acid PET or MR spectroscopy are employed [27]. In a prospective study of 35 patients with proven glioma, Karunanithi et al. found the sensitivity of ¹⁸F-FDOPA-PET/CT in determining recurrence in glioma to be 100% when compared to 92% with contrast-enhanced MRI and a specificity of 89% with ¹⁸F-FDOPA-PET/CT versus 44% for contrast enhanced-MRI [28]. Karunanithi et al. in a separate study of 28 patients with proven glioma compared ¹⁸F-FDOPA-PET/CT with ¹⁸F-Fluoro-deoxy-glucose (FDG)-PET/CT and found the sensitivity and specificity for FDG was inferior at 48% and 100% respectively and in comparison, ¹⁸F-FDOPA-PET/CT was 100% and 86% respectively [29]. A meta-analysis by de Zwart et al. found ¹⁸F-FDOPA to be superior over ¹¹C-MET and ¹⁸F-FET in differentiating tumour progression from treatment-related changes with a pooled sensitivity of 85-100% and specificity of 72-100% when compared to ¹¹C-MET (sensitivity 80–98%, specificity 61–91%) and ¹⁸F-FET (sensitivity 81–95%, specificity 71–93%) [30]. A study by Villani et al. with 50 patients found a potential role of FDOPA in prognostication of low-grade gliomas [31]. The authors found that disease duration and a maximum standardised uptake value (SUV_{max}) of > 1.75 was predictive of progression and superior to MRI in detection of progression and therefore prognosis, as transformation to high-grade is the key determinant in patient survival. When the recommendations were published there were conflicting results on the ability of ¹⁸F-FDOPA-PET to predict glioma grade and ¹⁸F-FDOPA was not recommended for this use. This article will review previously published results and results from recent studies that have become available since the publication of these guidelines.

¹⁸F-FDOPA PET

[¹⁸F]fluorodopa, 3, 4-dihydroxy-6-[¹⁸F]fluoro-L-phenylalanine. (¹⁸F-FDOPA) was originally developed for brain imaging in patients with movement disorders. It consists of an amino acid, phenylalanine, attached to a radioisotope, fluorine (Fig. 2a), that is able to cross the blood brain barrier and act as a precursor for dopamine. Phenylalanine is an essential aromatic amino acid with a neutral charge. Gliomas require a continuous supply of amino acids to maintain protein synthesis and cell proliferation. These amino acids reach the tumour cells via amino acid transporters. Amino acids are cationic, anionic or neutral, and their transport across a membrane is regulated by amino acid transporters [32]. Amino acid transporters can be uniporter, antiporter or symporter, each transporting certain amino acids (Fig. 2b). One of these neutral transporter systems is the L-type amino acid transporter (LAT) which is a membrane bound Na⁺ independent transport protein regulating the transport of essential amino acids across cell membranes. There are four main types of LAT transporter; LAT1 (SLC7A5), LAT2 (SLC7A8), LAT3 (SLC43A1) and LAT4 (SLC43A2). LAT1 is a polypeptide consisting of 507 amino acids and 12



Fig.2 a Chemical structure of $[^{18}F]$ fluorodopa (-NH₂=amine group, -COOH=carboxyl group). b Amino acids reach tumour cells via amino acid transporters. Amino acid transporters can be uniporter, symporter or antiporter and each have carrier numbers as demonstrated

transmembrane regions with a molecular weight of 55 kDa [33]. LAT1 forms a heterodimeric complex with CD98 via a disulphide bond. CD98 is a polypeptide of 630 amino acids and a molecular weight of 68 kDa. CD98 is thought to be crucial for LAT1 to function but it has been demonstrated that LAT1 is the only transport component of the LAT1/ CD98 heterodimer [33, 34]. Substrates that bind the LAT1 transporter must have a histidine group, a carboxylic group and an amino group [35]. LAT1 controls the movement of neutral essential amino acids which include histidine, leucine, isoleucine, methionine, phenylalanine, tyrosine, tryptophan and valine, into cells in exchange for the efflux of intracellular substrates such as glutamine, histidine and tyrosine (Fig. 3) [36, 37]. LAT1 mRNA is expressed most strongly in the brain, placenta, colon, testis and spleen and is expressed at low levels in the lungs, liver and heart [38]. Studies have demonstrated that LAT1 expression is higher than in normal tissues in cancers [39]. Tumour progression in gliomas requires a constant supply of amino acids for protein synthesis and cell proliferation. LAT1 expression correlates with proliferation of cancer cells enabling rapid growth as it plays role in cell growth, transcription and translation through the mammalian target of rapamycin (mTOR) signalling pathway which facilitates protein synthesis [35, 40].

When administered intravenously, 1% of the ¹⁸F-FDOPA will cross the blood brain barrier via the LAT1 transporter and the rest remains in the periphery where it is converted to 3-O-methyl-6-fluoro-L-DOPA (OMFD) by catechol O-methyl transferase (COMT) or into [¹⁸F]fluorodopamine by aromatic amino acid decarboxylase (AAAD). Once across the BBB FDOPA is converted to [¹⁸F]fluorodopamine by aromatic amino acid decarboxylase (AAAD) (Fig. 4). Fluorodopamine behaves like dopamine in vivo and



Fig. 3 Structure and function of the LAT1 amino acid transporter



Fig.4 Metabolism of FDOPA (AAAD=aromatic amino acid decarboxylase, FDA=18F6-dlurodopamine, COMT=catechol O-methyl transferase, OMFD=3-O-methyl-6-fluoro-L-DOPA, FDO-

 $PAC = [^{18}F]6$ -fluoro-L-3,4-dihydrophenylacetic acid, MAO = monoamine oxidase (MAO), FHVA = [^{18}F]6-fluorohomovanillic acid)

is either stored in pre-synaptic vesicles in the striatum or metabolised into [¹⁸F]6-fluoro-L-3,4-dihydrophenylacetic acid (FDOPAC) by monoamine oxidase (MAO) and then into [¹⁸F]6-fluorohomovanillic acid (FHVA) by COMT [41]. The ¹⁸F-FDOPA metabolites are renally excreted. In glioma cells the ¹⁸F-FDOPA is not metabolised [42]. To reduce the systemic metabolism of ¹⁸F-FDOPA and increase bio-availability and cerebral uptake, the decarboxylase inhibitor carbidopa is often administered prior to administration of ¹⁸F-FDOPA. Bros et al. found that in imaging of gliomas with ¹⁸F-FDOPA-PET, premedication with carbidopa resulted in a 50% increase in uptake in all brain structures but when corrected for the tumour-to-healthy-brain ratio did not impact image interpretation [43].

¹⁸F-FDOPA has been demonstrated to be advantageous over other amino acid PET tracers as it is predominantly transported by the L-type amino acid transporter without significant uptake into surrounding normal brain parenchyma with the exception of the basal ganglia, thereby allowing easier discrimination of uptake within the tumour [44, 45]. Despite this limitation, in gliomas involving the basal ganglia ¹⁸F-FDOPA has been shown be able to accurately delineate tumour boundaries [46, 47]. ¹⁸F-FDOPA is more readily available in clinical practice compared to ¹¹C-MET which requires an on-site cyclotron due to its short half-life of 20 minutes whereas the half-life of ¹⁸F-FDOPA is 110 minutes [48]. ¹⁸FDG has been extensively used for imaging in brain tumours with a meta-analysis of 26 studies that reported a sensitivity of 77% and specificity of 78% in detecting gliomas irrespective of grade [49–51]. However, ¹⁸FDG measures glucose metabolism and therefore it tends to accumulate in the grey matter which can interfere with the ability to differentiate tumour grade reliably. Tumours with low glucose metabolism, such as low-grade gliomas, are often not well visualised with FDG [52]. ¹⁸F-FDOPA has been shown to have a higher uptake in low-grade gliomas than FDG [53]. ¹⁸F-FDOPA uptake into glioma cells is thought to be higher in areas with high-grade features due to an increase in the transport of amino acids into tumour cells which is led by an increase in expression of the L-type amino acid transport system and subsequently has the potential to be able to detect areas of high-grade transformation within low-grade glioma before the typical radiological features of transformation such as contrast enhancement are visible on conventional MRI [54, 55]. Ledezma et al. found in a small number of cases that ¹⁸F-FDOPA tracer activity was able to identify tumour not visible on conventional MRI [52].

Role of ¹⁸F-FDOPA in differentiating highand low-grade gliomas

The standardised uptake value (SUV) is a measure of ¹⁸F-FDOPA uptake and is a calculation of the ratio of tissue radioactivity concentration (in kBq/ml) at a given time divided by the administered activity at the time of injection (in MBq) divided by the body weight (in kg) [56]. Despite studies investigating ¹⁸F-FDOPA SUV in differing grades of glioma there are currently no agreed thresholds for SUV in routine clinical practice with ¹⁸F-FDOPA for distinguishing between high and low-grade gliomas. The Joint EANM/EANO/RANO guidelines do however provide thresholds for ¹⁸F-FDOPA in delineating tumour extent, detecting tumour recurrence and identification of response to treatment with bevacizumab [18]. For extent of tumour Pafundi et al. identified that a tumour-to-normal-brain ratio (TBR) of greater than 2.0 corresponded to high-grade disease [57]. For detection of tumour recurrence, a tumour to striatum ratio (TSR) max of 2.1 and a TSR_{mean} of 1.8 are described [58]. Scwharzenberg et al. were able to show that by using ¹⁸F-FDOPA-PET at two weeks following initiation of bevacizumab therapy for recurrent high-grade gliomas the threshold for a positive response to treatment was found to be a brain tumour volume decrease of greater than 35% or a metabolically active tumour volume of less than 18 mL at two weeks [24].

Literature published prior to joint EANM/EANO/ RANO guidelines

Prior to the publication of the 2016 recommendations there were two main studies that have addressed ¹⁸F-FDOPA PET in predicting glioma grade. The first was by Pafundi et al. who performed a prospective pilot study with 10 patients [57]. Of these, 8 were newly diagnosed and 2 were recurrent gliomas. For the patients undergoing surgical resection, a maximum of three stereotactic biopsy targets were planned using various PET SUVs along a single trajectory with the PET/CT and contrast enhanced MRI fused using MIM Maestro software. 23 biopsy samples were obtained using neuronavigation on the preplanned targets and each tissue sample was analysed and graded along with recording the average cellularity and average Ki-67 index. PET/CT was performed 10 min after the tracer injection and carbidopa premedication was not used. A strong association was found of ¹⁸F-FDOPA SUV_{max} with tumour grade. The authors identified that there were significant differences between distinguishing grade II and IV (p=0.008) and grade III and IV (p=0.024) when using static ¹⁸F-FDOPA-PET/CT in 8 patients. No significant difference was found between the SUV_{max} in grade II and grade III tumours (p=0.17). This may have been because there were only two grade II astrocytomas in the cohort. By removing the oligodendroglioma samples the authors found a significant correlation between ¹⁸F-FDOPA SUV_{mean} and histological cellularity (p=0.01). Higher cellularity indicates histologically higher-grade features. A TBR of > 2.0 was able to define high-grade components of astrocytic tumours. The oligodendroglioma biopsy samples, of which there were three, were removed from the final analysis as the study found that the ¹⁸F-FDOPA SUV_{max} was much greater in comparison to the grade II astrocytomas and this has been previously described with the tracers 11 C-MET⁵⁵ and 18 F-FET [60].

The second study by Fueger et al. included 59 patients of which 22 where newly diagnosed and 37 recurrent gliomas of varying grades (grade II n = 22, grade III n = 19, grade IV n = 27) that underwent static ¹⁸F-FDOPA-PET/CT before

surgery [61]. The PET/CT was performed 10 min after the tracer injection and carbidopa premedication was not administered. Each tumour was graded histologically and Ki-67 expression measured. The authors found a significant difference in SUV_{max} in newly diagnosed gliomas between grade II and III (p=0.044), between grade II and IV (p=0.007)and between grade III and IV tumours (p=0.010) and concluded that a ¹⁸F-FDOPA SUV_{max} of 2.72 was the cut-off to distinguish low- and high-grade newly diagnosed gliomas. There was no significant difference between grades in the recurrent tumours. The lack of correlation in tumour recurrence could be explained by damage to the BBB from radiation therapy leading to increased vascular permeability resulting in an increase in non-carrier (LAT1) mediated transport of ¹⁸F-FDOPA from endothelial cells into tumour cells [62]. The histopathological samples from this study were much larger compared to Pafundi et al. [57] and therefore would have not considered ¹⁸F-FDOPA uptake heterogeneity within the tumour.

Literature published after Joint EANM/EANO/RANO guidelines available

The following section will describe the new data published following the 2016 recommendations and the impact of this data. Todeschi et al. recently performed a single centre prospective study on 16 newly diagnosed gliomas and 4 recurrent gliomas using static ¹⁸F-FDOPA-PET/CT [63]. The PET/CT was performed 30 min after tracer injection and no carbidopa premedication was administered. Biopsy targets were based on regions of tracer hypermetabolism and hypometabolism. In this series the authors found that a SUV_{max} threshold of > 1.75 demonstrated a greater yield in terms of diagnosis of high-grade gliomas. The authors concluded that the use of ¹⁸F-FDOPA-PET allowed for better targeting of metabolically active areas of tumour to reduce biopsy sampling bias but when used alone was unable to reliably distinguish tumour grade (SUV $_{max}$ low-grade 2.03 and 2.18 for high-grade, p = 0.64). The biopsies were taken using a robotic arm with a biopsy needle as opposed to a craniotomy for tumour resection as in the Pafundi study [57]. The result is that the preplanned targets and resultant biopsies are likely to have been more accurate as they would have not been affected by brain shift from opening the dura and CSF drainage as would have been the case with the Pafundi study [57]. However, the single trajectory resulted in two biopsy targets (an area of hypometabolism and hypermetabolism) that were in close proximity with little margin for error in coregistration of the patient with the imaging.

The majority of ¹⁸F-FDOPA-PET studies have used static parameters for ¹⁸F-FDOPA uptake. Static PET provides a single snapshot of the tracer uptake whereas dynamic PET takes multiple snapshots at different timepoints. The advantage of dynamic PET when used alongside kinetic modelling allows creation of time-activity curves (TAC) which can provide additional information on the pharmacokinetics of ¹⁸F-FDOPA in brain tumours. The assumption is that high-grade gliomas will quickly take up ¹⁸F-FDOPA (the wash-in) when injected in comparison to low-grade gliomas which will have a slow wash-in period [64]. Previously Schiepers et al. have directly compared static and dynamic ¹⁸F-FDOPA-PET in brain tumours [42]. In this study 37 patients were included of which 33 were primary brain tumours. A significant difference was found between tracer volume distribution between newly diagnosed lowgrade and high-grade tumours (p < 0.01) and between newly diagnosed high-grade tumours and tumours with post-treatment changes. Nioche et al. similarly performed static and dynamic ¹⁸F-FDOPA-PET/CT for 33 patients published in 2013 and found a SUV_{mean} threshold of 2.5 in determining the grade with a sensitivity of 94% and specificity of 66% [65]. There was no significant improvement in these results when comparing static and dynamic imaging. Dynamic ¹⁸F-FDOPA-PET appears to be more useful in newly diagnosed gliomas and less useful for detection of recurrence or progression as evidence by Zaragori et al. [66]. This study identified 51 patients with suspected glioma recurrence or progression who underwent ¹⁸F-FDOPA-PET and found that no additional significant information was gained from performing dynamic imaging. Xiao et al. in a recent metaanalysis reported a pooled sensitivity of 0.71 and specificity of 0.86 for grading newly diagnosed gliomas with static FDOPA PET [67].

Despite the sensitivity and specificity of static ¹⁸F-FDOPA-PET, dynamic ¹⁸F-FDOPA-PET has the advantage of being able to detect a tumour's molecular characteristics as demonstrated by Ginet et al. [68]. This retrospective study looked at 58 patients with newly diagnosed glioma who underwent either biopsy (n = 24) or surgical resection (n=34) and preoperative static and dynamic ¹⁸F-FDOPA-PET. Patients were given carbidopa one hour prior to PET imaging. They found that only the dynamic parameters of time-to-peak (TTP) which represents the time from tracer injection to maximum SUV, area under the curve (AUC) and curve slopes were significant in predicting the IDH-mutation status (TTP $p \le 0.001$, AUC p = 0.789, slope p = 0.013). For prediction of the 1p/19q co-deletion status the TTP was the only dynamic parameter found to be statistically significant (p=0.034). The static imaging did not significantly correlate with the molecular subtypes. This could be because static imaging is just a snapshot in time whereas dynamic imaging captures the rate of ¹⁸F-FDOPA tracer uptake and washout so more information is available. Isal et al. retrospectively looked at 20 patients with histologically confirmed newly diagnosed grade II and grade III gliomas that had serial static ¹⁸F-FDOPA PET [69]. The authors found that a SUV_{max} of greater than 1.8 is predictive of the presence of an IDH-mutation. Similarly, Cicone et al. performed static ¹⁸F-FDOPA-PET/CT in 33 patients following surgery but before initiation of chemoradiotherapy [70]. The SUV_{max}, TBR and TSR were not statistically significant between tumours that were IDH-mutant or IDH-wildtype and that no difference was found between 1p/19q co-deleted and nonco-deleted patients. In addition, no statistically significant difference was seen between low- and high-grade gliomas ($p \ge 0.2$).

The presence of MGMT methylation indicates silencing of the MGMT gene resulting in a reduction in the capability of tumour cells to repair damage from alkylating agents such as temozolomide and confers a better prognosis [71]. Cimini et al. performed static ¹⁸F-FDOPA-PET/CT in 72 patients post-surgery and found no difference between the presence of MGMT methylation versus unmethylation (p=0.15) and no difference between IDH-mutant and IDH-wildtype gliomas (p=0.79) [72]. One explanation for the results in both these studies may be that the ¹⁸F-FDOPA-PET scans were performed following surgical biopsy and it has already been shown that inflammation and macrophage response seen post-surgery can alter ¹⁸F-FDOPA uptake [73, 74]. The ability to detect IDH-mutation status on imaging is novel and could potentially play a role in management discussions with patient's at diagnosis as IDH-wildtype tumours regardless of grade have a shorter median overall survival, <2 years, when compared to IDH-mutant gliomas [75].

The GLIROPA clinical trial form Girard et al. included only newly diagnosed diffuse gliomas of which 32 biopsies were acquired from 14 patients [56]. Preoperative static and dynamic ¹⁸F-FDOPA-PET was performed followed by stereotactically guided biopsies (up to three per patient). The PET/CT was performed 10 min after tracer injection and no carbidopa premedication was administered. Each biopsy sample was graded independently but there was a much higher number of high-grade tumour samples (n=23)in comparison to low-grade tumour samples (n=9). The authors established that the static PET parameters were not significantly different between grades but the kinetic analysis from dynamic image acquisition was more accurate for glioma grading. However, the time between the ¹⁸F-FDOPA-PET/CT and stereotactic biopsy to confirm the histopathological grade was as long as 110 days during which time the tumour had the potential to transform to a higher grade. Janvier et al. performed a retrospective review published in 2015 on 31 patients of which 6 were recurrent gliomas and 25 newly diagnosed who had undergone static ¹⁸F-FDOPA-PET [76]. The study found that the SUV_{mean} and tumourto-normal-tissue ratio (T/N) best correlated to the grade (p = < 0.05) with a cut-off for SUV_{mean} of 1.33. Bund et al. found that in a subset of low-grade gliomas, in discriminating between dysembryoplastic neuroepithelial tumour

 Table 1
 Summary of spatially linked ¹⁸F-FDOPA-PET studies

Authors	Year	Num- ber of patients	Number of newly diagnosed gliomas	Number of recurrent gliomas	Type of imaging	Number of biopsies per- formed	Type of study	FDOPA thresholds
Girard et al	2021	14	14	0	Dynamic PET/CT	32	Prospective	None given
Pafundi et al	2013	10	8	2	Static PET/CT	23	Prospective	TBR > 2.0 equates to high-grade components of astrocytic tumours
Ponisio et al	2020	10	6	4	Dynamic PET/ MRI	23	Prospective	None given
Todeschi et al	2019	20	16	4	Static PET/CT	20	Prospective	SUVmax > 1.75 gave a higher yield for high grade

and grade II oligodendroglioma SUV_{max} was significant ($p \le 0.01$) and also between low-and high-grade gliomas a SUV_{max} cut off of 2.16 was found [77]. A prospective study in 45 patients with suspected glioma who underwent preintervention static ¹⁸F-FDOPA-PET and found that a T/N SUV_{max} ratio greater 1.7 was able to differentiate high-grade glioma from other graded lesions [78].

In a pilot study Ponisio et al. used dynamic ¹⁸F-FDOPA-PET/MRI with stereotactically linked histopathology data in 10 patients of which 4 were recurrent tumours, obtaining a total of 23 biopsies. The authors identified that the results of the ¹⁸F-FDOPA-PET/MRI had a positive impact on patient management in 4 cases including performing additional biopsies and altering surgical strategies in terms of extent of resection. In addition, the authors found a strong correlation between tumour SUV parameters and the Ki-67 index reflecting cell proliferation. Despite the promising results and impact on management Ponisio et al. concluded that dynamic ¹⁸F-FDOPA-PET/MRI was independent of WHO grade and did not significantly differentiate between low- and high-grade gliomas. This pilot study and Todeschi et al. [63] differs from the other studies with stereotactically linked histopathological data [56, 57] in that the ¹⁸F-FDOPA-PET was combined with MRI rather than CT. There are implications in terms of cost, reduction in radiation exposure and accessibility in using PET/MRI but they are reported to perform equally [79].

There has been a lot of work on the ability of ¹⁸F-FDOPA-PET to differentiate between low- and high-grade gliomas since the 2016 recommendations were published which have been discussed but there are a number of limitations to consider. Firstly, many studies include both newly diagnosed and recurrent gliomas together. Macrophages have been reported to have high levels of amino acid transport and therefore ¹⁸F-FDOPA uptake. As a result, ¹⁸F-FDOPA uptake levels may be falsely positive following surgery and radiotherapy due to the presence of macrophages around the resection cavity and in irradiated tumours [73, 74]. For this reason, ideally newly diagnosed and recurrent gliomas should be reviewed separately to exclude treatment bias and if not possible, the time delay between surgery and ¹⁸F-FDOPA imaging should be considered and any uptake around a resection cavity should be interpreted with care. Secondly, in the majority of studies the number of participants is generally low and there are differing protocols from the biopsy and surgical technique to the co-registration of PET/CT with MRI and histological interpretation of tissue samples and therefore it is impossible to completely merge data from different studies. In addition, there are few studies [56, 57, 63, 80] that have integrated biopsy location with histopathological data and ¹⁸F-FDOPA uptake. This is important as ¹⁸F-FDOPA uptake can vary widely within a tumour. The ¹⁸F-FDOPA uptake heterogeneity may represent heterogeneity in terms of the histopathological features and therefore grade, which can alter management strategies and ultimately prognosis. Finally, the current published studies are all single centre and there is a requirement here for multicentre studies to be completed.

Future work

Additional larger prospective studies are needed with biopsy validated dynamic ¹⁸F-FDOPA-PET/CT and PET/MRI in treatment naïve gliomas to confirm the promising correlation seen in the literature. A stage 2 clinical trial run by the Mayo Clinic (Rochester, United States) has now closed to enrolment having recruited 72 patients. With the aim of determining ¹⁸F-FDOPA-PET thresholds for distinguishing low- from high-grade glioma [81]. The FIG (¹⁸F-FDOPA-PET PET imaging in glioma) study is currently recruiting patients in Oxford University Hospitals NHS Foundation

Authors	Year	Total number of patients	Number of newly diagnosed gliomas	Num- ber of recurrent gliomas	Type of imaging used	Type of study	Key Findings
Fueger et al	2010	59	22	37	Static ¹⁸ F-FDOPA- PET/CT	Combination of prospective + retro- spective	¹⁸ F-FDOPA SUV _{max} cut-off of 2.72 to dis- tinguish low- and high- grade newly diagnosed gliomas
							No significant difference in recurrent tumours
Schiepers et al	2007	37	33	4	Static and dynamic ¹⁸ F-FDOPA-PET/CT	Prospective	Significant difference between tracer volume distribution of newly diagnosed low + high- grade tumours $(p \le 0.01)$
Nioche et al	2013	33	20	13	Static and dynamic ¹⁸ F-FDOPA-PET/CT	Prospective	SUV _{mean} threshold of 2.5 in determining the grade (sensitivity 94%, specificity 66%)
							No significant improve- ment when compar- ing static + dynamic imaging
Zaragori et al	2020	51	0	51	Dynamic ¹⁸ F-FDOPA- PET/CT	Retrospective	No additional infor- mation gained from dynamic imaging
Ginet et al	2020	58	58	0	Static and dynamic ¹⁸ F-FDOPA-PET/CT	Retrospective	TTP and AUC sig- nificant in predicting IDH-mutation status (TTP $p \le 0.001$, AUC p = 0.789, slope p = 0.013)
							TTP significant in prediction of 1p/19q co-deletion status (p=0.034)
Isal et al	2018	20	20	0	Static ¹⁸ F-FDOPA- PET/CT	Retrospective	SUV _{max} > 1.8 predictive of the presence of an IDH-mutation
Cicone et al	2019	33	33	0	Static ¹⁸ F-FDOPA- PET/CT	Prospective	No significant differ- ence seen between low-+high-grade gliomas ($p \ge 0.2$)
Cimini et al	2020	72	Not commented on	Not com- mented on	Static ¹⁸ F-FDOPA- PET/CT	Retrospective	No difference between presence of MGMT methylation ver- sus unmethylation (p=0.15)
							No difference between IDH-mutant and IDH-wildtype gliomas (p=0.79)
Janvier et al	2015	31	25	6	Static ¹⁸ F-FDOPA- PET/CT	Retrospective	SUV_{mean} + T/N ratio correlated to grade (p = < 0.05)

Table 2 Summary of non-spatially linked ¹⁸F-FDOPA-PET studies and the key findings

 Table 2 (continued)

Authors	Year	Total number of patients	Number of newly diagnosed gliomas	Num- ber of recurrent gliomas	Type of imaging used	Type of study	Key Findings
Bund et al	2017	53	53	0	Static ¹⁸ F-FDOPA- PET/CT	Prospective	SUV _{max} cut off 2.16 sig- nificant for discriminat- ing low-+high-grade glioma
Patel et al	2018	45	45	0	Static ¹⁸ F-FDOPA- PET/CT	Retrospective	T/N SUV _{max} ratio > 1.7 able to differentiate high-grade glioma from other graded lesions

Trust (Oxford, UK) with the aim of opening to recruitment at two further sites later this year. The FIG study is a feasibility study investigating ¹⁸F-FDOPA-PET and oxygen-enhanced MR guided histopathology in patients with suspected low-grade gliomas in a multi-centre setting [82]. The study will include up to 168 biopsies in 21 participants recruited from multiple neurosurgical centres in the UK. This study will be one of the first and largest study to use dynamic FDOPA PET/CT in determining grade of gliomas in a multi-centre setting.

The recent publication of the 2021 WHO classification of central nervous system tumours has incorporated numerous molecular changes that support the integrated diagnosis for each tumour type¹ (Table 1). These additional molecular changes open up an opportunity for evaluation of dynamic ¹⁸F-FDOPA-PET in detection of molecular markers similar to the previous studies into IDH-mutation status, 1p/19q codeletion status and presence of MGMT methylation [68–72] (See Table 2).

Conclusions

The literature demonstrates the capability of ¹⁸F-FDOPA-PET to differentiate low- from high-grade gliomas as well as the presence of an IDH-mutation, the 1p/19q co-deletion status and the MGMT methylation status. The FIG multicentre study will possibly answer questions on the introduction of preoperative ¹⁸F-FDOPA-PET into the routine imaging work up for glioma patients to guide surgical management. Detecting areas of high-grade transformation more effectively would allow for earlier clinical intervention, optimisation of surgical planning and resection strategies with the potential to lengthen progression free survival and therefore ultimately prognosis. Author contributions All authors contributed to the study conception and design. Material preparation, data collection and analysis were performed by Joy Roach. The first draft of the manuscript was written by Joy Roach and all authors commented on previous versions of the manuscript. All authors read and approved the final manuscript.

Funding This work was funded by the CRUK National Cancer Imaging Translational Accelerator (C42780/A27066).

Declarations

Conflict of interest The authors have not disclosed any competing interests.

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit http://creativecommons.org/licenses/by/4.0/.

References

- Louis DN, Perry A, Wesseling P et al (2021) The WHO classification of tumors of the central nervous system: a summary. Neuro Oncol 23(8):1231–1251. https://doi.org/10.1093/neuonc/noab106
- Smits M (2016) Imaging of oligodendroglioma. Br J Radiol 89(1060):20150857. https://doi.org/10.1259/bjr.20150857
- Henson JW, Gaviani P, Gonzalez RG (2005) MRI in treatment of adult gliomas. Lancet Oncol 6(3):167–175. https://doi.org/10. 1016/S1470-2045(05)01767-5
- Muragaki Y, Chernov M, Maruyama T et al (2008) Low-grade glioma on stereotactic biopsy: how often is the diagnosis accurate? Minim Invasive Neurosurg 51(5):275–279. https://doi.org/ 10.1055/s-0028-1082322
- Jackson RJ, Fuller GN, Abi-Said D et al (2001) Limitations of stereotactic biopsy in the initial management of gliomas. Neuro-Oncology 3(3):193-200

- Beiko J, Suki D, Hess KR et al (2014) IDH1 mutant malignant astrocytomas are more amenable to surgical resection and have a survival benefit associated with maximal surgical resection. Neuro Oncol 16(1):81–91
- Kim BYS, Jiang W, Beiko J et al (2014) Diagnostic discrepancies in malignant astrocytoma due to limited small pathological tumor sample can be overcome by IDH1 testing. J Neurooncol 118:405–412
- Padelli F, Mazzi F, Erbetta A, et al (2022) In vivo brain MR spectroscopy in gliomas: clinical and pre-clinical chances. Clinical and Translational Imaging 10:495–515
- 9. Hilario A, Ramos A, Perez-Nuñez A et al (2012) The added value of apparent diffusion coefficient to cerebral blood volume in the preoperative grading of diffuse gliomas. AJNR Am J Neuroradiol 33(4):701–707. https://doi.org/10.3174/ajnr.A2846
- Law M, Yang S, Wang H et al (2003) Glioma grading: sensitivity, specificity, and predictive values of perfusion MR imaging and proton MR spectroscopic imaging compared with conventional MR imaging. AJNR Am J Neuroradiol 24(10):1989–1998
- Toyooka M, Kimura H, Uematsu H, Kawamura Y, Takeuchi H, Itoh H (2008) Tissue characterization of glioma by proton magnetic resonance spectroscopy and perfusion-weighted magnetic resonance imaging: glioma grading and histological correlation. Clin Imaging 32(4):251–258. https://doi.org/10.1016/j.clinimag. 2007.12.006
- Palumbo B, Buresta T, Nuvoli S et al (2014) SPECT and PET serve as molecular imaging techniques and in vivo biomarkers for brain metastases. Int J Mol Sci 15(6):9878–9893. https://doi. org/10.3390/ijms15069878
- Chen W (2007) Clinical applications of PET in brain tumors. J Nucl Med 48(9):1468–1481. https://doi.org/10.2967/jnumed.106. 037689
- Galldiks N, Lohmann P, Cicone F, Langen K (2019) FET and FDOPA PET Imaging in Glioma. In: Pope W (ed) Glioma Imaging. Springer International Publishing, pp211–221
- 15 O'Connor JPB, Robinson SP, Waterton JC (2019) Imaging tumour hypoxia with oxygen-enhanced MRI and BOLD MRI. BJR. https://doi.org/10.1259/bjr.20180642
- National Institute for Health and Care Excellence. (2021). Brain tumours (primary) and brain metastases in over 16s [NICE Guideline No.99]. https://www.nice.org.uk/guidance/ng99
- Stupp R, Mason WP, van den Bent MJ et al (2005) Radiotherapy plus concomitant and adjuvant temozolomide for glioblastoma. N Engl J Med 352(10):987–996. https://doi.org/10.1056/NEJMo a043330
- Law I, Albert NL, Arbizu J et al (2019) Joint EANM/EANO/ RANO practice guidelines/SNMMI procedure standards for imaging of gliomas using PET with radiolabelled amino acids and [18F]FDG: version 1.0. Eur J Nucl Med Mol Imaging 46(3):540– 557. https://doi.org/10.1007/s00259-018-4207-9
- Albert NL, Weller M, Suchorska B et al (2016) Response assessment in neuro-oncology working group and european association for neuro-oncology recommendations for the clinical use of PET imaging in gliomas. Neuro Oncol 18(9):1199–1208. https://doi.org/10.1093/neuonc/now058
- Galldiks N, Langen KJ, Holy R et al (2012) Assessment of treatment response in patients with glioblastoma using O-(2–18Ffluoroethyl)-L-tyrosine PET in comparison to MRI. J Nucl Med 53(7):1048–1057. https://doi.org/10.2967/jnumed.111.098590
- Galldiks N, Kracht LW, Burghaus L et al (2006) Use of 11C-methionine PET to monitor the effects of temozolomide chemotherapy in malignant gliomas. Eur J Nucl Med Mol Imaging 33(5):516–524. https://doi.org/10.1007/s00259-005-0002-5
- 22. Jansen NL, Suchorska B, Schwarz SB et al (2013) [18F]fluoroethyltyrosine-positron emission tomography-based therapy

monitoring after stereotactic iodine-125 brachytherapy in patients with recurrent high-grade glioma. Mol Imaging 12(3):137-147

- 23. Harris RJ, Cloughesy TF, Pope WB et al (2012) 18F-FDOPA and 18F-FLT positron emission tomography parametric response maps predict response in recurrent malignant gliomas treated with bevacizumab. Neuro Oncol 14(8):1079–1089. https://doi.org/10. 1093/neuonc/nos141
- Schwarzenberg J, Czernin J, Cloughesy TF et al (2014) Treatment response evaluation using 18F-FDOPA PET in patients with recurrent malignant glioma on bevacizumab therapy. Clin Cancer Res 20(13):3550–3559. https://doi.org/10.1158/1078-0432. CCR-13-1440
- Langen KJ, Galldiks N, Hattingen E, Shah NJ (2017) Advances in neuro-oncology imaging. Nat Rev Neurol 13(5):279–289. https:// doi.org/10.1038/nrneurol.2017.44
- 26 Galldiks N, Lohmann P, Albert NL, Tonn JC, Langen KJ (2019) Current status of PET imaging in neuro-oncology. Neurooncol Adv. https://doi.org/10.1093/noajnl/vdz010
- Ellingson BM, Chung C, Pope WB, Boxerman JL, Kaufmann TJ (2017) Pseudoprogression, radionecrosis, inflammation or true tumor progression? challenges associated with glioblastoma response assessment in an evolving therapeutic landscape. J Neurooncol 134(3):495–504. https://doi.org/10.1007/ s11060-017-2375-2
- Karunanithi S, Sharma P, Kumar A et al (2013) Comparative diagnostic accuracy of contrast-enhanced MRI and (18)F-FDOPA PET-CT in recurrent glioma. Eur Radiol 23(9):2628–2635. https:// doi.org/10.1007/s00330-013-2838-6
- Karunanithi S, Sharma P, Kumar A et al (2013) 18F-FDOPA PET/ CT for detection of recurrence in patients with glioma: prospective comparison with 18F-FDG PET/CT. Eur J Nucl Med Mol Imaging 40(7):1025–1035. https://doi.org/10.1007/s00259-013-2384-0
- 30. de Zwart PL, van Dijken BRJ, Holtman GA, Stormezand GN, Dierckx RAJO, Jan van Laar P et al (2020) Diagnostic accuracy of PET tracers for the differentiation of tumor progression from treatment-related changes in high-grade glioma: a systematic review and metaanalysis. J Nucl Med 61(4):498–504
- Villani V, Carapella CM, Chiaravalloti A et al (2015) The role of PET [18F]FDOPA in evaluating low-grade Glioma. Anticancer Res 35(9):5117–5122
- 32. Gauthier-Coles G, Vennitti J, Zhang Z et al (2021) Quantitative modelling of amino acid transport and homeostasis in mammalian cells. Nat Commun 12(1):5282. https://doi.org/10.1038/ s41467-021-25563-x
- Napolitano L, Scalise M, Galluccio M, Pochini L, Albanese LM, Indiveri C (2015) LAT1 is the transport competent unit of the LAT1/CD98 heterodimeric amino acid transporter. Int J Biochem Cell Biol 67:25–33. https://doi.org/10.1016/j.biocel.2015.08.004
- 34 Puris E, Gynther M, Auriola S, Huttunen KM (2020) L-Type amino acid transporter 1 as a target for drug delivery. Pharm Res 37(5):88. https://doi.org/10.1007/s11095-020-02826-8
- 35. Scalise M, Galluccio M, Console L, Pochini L, Indiveri C (2018) The human SLC7A5 (LAT1): the intriguing histidine/large neutral amino acid transporter and its relevance to human health. Front Chem 6:243. https://doi.org/10.3389/fchem.2018.00243
- Zaragozá R (2020) Transport of amino acids across the bloodbrain barrier. Front Physiol 11:973. https://doi.org/10.3389/fphys. 2020.00973
- Herholz K (2017) Brain tumors: an update on clinical PET research in gliomas. Semin Nucl Med 47(1):5–17. https://doi.org/ 10.1053/j.semnuclmed.2016.09.004
- Kanai Y, Segawa H, Miyamoto K-i, Uchino H, Takeda E, Endou H (1998) Expression cloning and characterization of a transporter for large neutral amino acids activated by the heavy chain of 4F2 Antigen (CD98)*. J Biol Chem 273(37):23629–23632

- Zhang J, Xu Y, Li D et al (2020) Review of the correlation of LAT1 with diseases: mechanism and treatment. Front Chem 8:564809. https://doi.org/10.3389/fchem.2020.564809
- Wang Q, Holst J. 2015 L-type amino acid transport and cancer: targeting the mTORC1 pathway to inhibit neoplasia. American Journal of Cancer Research. 5(4):1281-94
- Endres CJ, DeJesus OT, Uno H, Doudet DJ, Nickles JR, Holden JE (2004) Time profile of cerebral [18F]6-fluoro-L-DOPA metabolites in nonhuman primate: implications for the kinetics of therapeutic L-DOPA. Front Biosci 9:505–512. https://doi.org/10.2741/ 1224
- 42 Schiepers C, Chen W, Cloughesy T, Dahlbom M, Huang S-C (2007) 18F-FDOPA kinetics in brain tumors. J Nuclear Med. https://doi.org/10.2967/jnumed.106.039321
- 43 Bros M, Zaragori T, Rech F et al (2021) Effects of carbidopa premedication on 18F-FDOPA PET imaging of glioma: a multiparametric Analysis. Cancers (Basel). https://doi.org/10.3390/ cancers13215340
- 44. Ono M, Oka S, Okudaira H et al (2013) Comparative evaluation of transport mechanisms of trans-1-amino-3-[(1)(8)F]fluorocyclobutanecarboxylic acid and L-[methyl-(1)(1)C]methionine in human glioma cell lines. Brain Res 1535:24–37. https://doi.org/10.1016/j. brainres.2013.08.037
- 45. Oka S, Okudaira H, Ono M et al (2014) Differences in transport mechanisms of trans-1-amino-3-[18F]fluorocyclobutanecarboxylic acid in inflammation, prostate cancer, and glioma cells: comparison with L-[methyl-11C]methionine and 2-deoxy-2-[18F] fluoro-D-glucose. Mol Imaging Biol 16(3):322–329. https://doi. org/10.1007/s11307-013-0693-0
- 46. Becherer A, Karanikas G, Szabó M et al (2003) Brain tumour imaging with PET: a comparison between [18F]fluorodopa and [11C]methionine. Eur J Nucl Med Mol Imaging 30(11):1561– 1567. https://doi.org/10.1007/s00259-003-1259-1
- Morana G, Puntoni M, Garrè ML, Massollo M, Lopci E, Naseri M et al (2016) Ability of (18)F-DOPA PET/CT and fused (18) F-DOPA PET/MRI to assess striatal involvement in paediatric glioma. Eur J Nucl Med Mol Imaging 43(9):1664–1672
- Lapa C, Linsenmann T, Monoranu CM, Samnick S, Buck AK, Bluemel C et al (2014) Comparison of the amino acid tracers 18F-FET and 18F-DOPA in high-grade glioma patients. J Nucl Med 55(10):1611–1616
- Nihashi T, Dababreh I, Terasawa T (2013) Diagnostic accuracy of PET for recurrent glioma diagnsois: a meta-analysis. AJNR 34(5):941–1011
- 50 Nihashi T, Dahabreh IJ, Terasawa T (2013) Diagnostic accuracy of PET for recurrent glioma diagnosis: a meta-analysis. AJNR Am J Neuroradiol 34(5):944–950. https://doi.org/10.3174/ajnr.A3324
- 51. O'Connor JP, Naish JH, Parker GJ et al (2009) Preliminary study of oxygen-enhanced longitudinal relaxation in MRI: a potential novel biomarker of oxygenation changes in solid tumors. Int J Radiat Oncol Biol Phys 75(4):1209–1215. https://doi.org/10. 1016/j.ijrobp.2008.12.040
- 52. Ledezma CJ, Chen W, Sai V et al (2009) 18F-FDOPA PET/MRI fusion in patients with primary/recurrent gliomas: initial experience. Eur J Radiol 71(2):242–248. https://doi.org/10.1016/j.ejrad. 2008.04.018
- Chen W, Silverman D, Delaloye S et al (2006) 18F-FDOPA PET imaging of brain tumors: comparison study with 18F-FDG PET and evaluation of diagnostic accuracy. J Nucl Med 47(6):904–911
- Youland RS, Kitange GJ, Peterson TE et al (2013) The role of LAT1 in 1^{8F}-DOPA uptake in malignant gliomas. J Neurooncol 111:11–18
- 55. Nawashiro H, Otani N, Uozumi Y et al (2005) High expression of L-type amino acid transporter 1 in infiltrating glioma cells. Brain Tumor Pathol 22(2):89–91. https://doi.org/10.1007/s10014-005-0188-z

- Girard A, Le Reste PJ, Metais A et al (2021) Additive value of dynamic FDOPA PET/CT for glioma grading. Front Med (Lausanne) 8:705996. https://doi.org/10.3389/fmed.2021.705996
- 57. Pafundi DH, Laack NN, Youland RS et al (2013) Biopsy validation of 18F-DOPA PET and biodistribution in gliomas for neurosurgical planning and radiotherapy target delineation: results of a prospective pilot study. Neuro Oncol 15(8):1058–1067. https:// doi.org/10.1093/neuonc/not002
- Herrmann K, Czernin J, Cloughesy T et al (2014) Comparison of visual and semiquantitative analysis of 18F-FDOPA-PET/CT for recurrence detection in glioblastoma patients. Neuro Oncol 16(4):603–609. https://doi.org/10.1093/neuonc/not166
- Hatakeyama T, Kawai N, Nishiyama Y et al (2008) 11C-methionine (MET) and 18F-fluorothymidine (FLT) PET in patients with newly diagnosed glioma. Eur J Nucl Med Mol Imaging 35(11):2009–2017. https://doi.org/10.1007/s00259-008-0847-5
- Jansen NL, Schwartz C, Graute V et al (2012) Prediction of oligodendroglial histology and LOH 1p/19q using dynamic [(18)F] FET-PET imaging in intracranial WHO grade II and III gliomas. Neuro Oncol 14(12):1473–1480. https://doi.org/10.1093/neuonc/ nos259
- Fueger BJ, Czernin J, Cloughesy T et al (2010) Correlation of 6–18F-fluoro-L-dopa PET uptake with proliferation and tumor grade in newly diagnosed and recurrent gliomas. J Nucl Med 51(10):1532–1538. https://doi.org/10.2967/jnumed.110.078592
- Langen KJ, Mühlensiepen H, Holschbach M, Hautzel H, Jansen P, Coenen HH (2000) Transport mechanisms of 3-[123I]iodo-alphamethyl-L-tyrosine in a human glioma cell line: comparison with [3H]methyl]-L-methionine. J Nucl Med 41(7):1250–1255
- Todeschi J, Bund C, Cebula H et al (2019) Diagnostic value of fusion of metabolic and structural images for stereotactic biopsy of brain tumors without enhancement after contrast medium injection. Neurochirurgie 65(6):357–364. https://doi.org/10.1016/j. neuchi.2019.08.002
- 64 Verger A, Imbert L, Zaragori T (2021) Dynamic amino-acid PET in neuro-oncology: a prognostic tool becomes essential. Eur J Nucl Med Mol Imaging. https://doi.org/10.1007/s00259-021-05530-w
- 65. Nioche C, Soret M, Gontier E et al (2013) Evaluation of quantitative criteria for glioma grading with static and dynamic 18F-FDopa PET/CT. Clin Nucl Med 38(2):81–87. https://doi. org/10.1097/RLU.0b013e318279fd5a
- 66. Zaragori T, Ginet M, Marie PY et al (2020) Use of static and dynamic [18F]-F-DOPA PET parameters for detecting patients with glioma recurrence or progression. EJNMMI Res 10(1):56. https://doi.org/10.1186/s13550-020-00645-x
- 67. Xiao J, Jin Y, Nie J, Chen F, Ma X (2019) Diagnostic and grading accuracy of 18FFDOPA PET and PET/CT in patients with gliomas: a systematic review and metaanalysis. BMC Cancer 19(1):767
- Ginet M, Zaragori T, Marie PY et al (2020) Integration of dynamic parameters in the analysis of 18F-FDopa PET imaging improves the prediction of molecular features of gliomas. Eur J Nucl Med Mol Imaging 47(6):1381–1390. https://doi.org/10. 1007/s00259-019-04509-y
- 69. Isal S, Gauchotte G, Rech F et al (2018) A high 18F-FDOPA uptake is associated with a slow growth rate in diffuse grade II-III gliomas. Br J Radiol 91(1084):20170803. https://doi.org/10.1259/ bjr.20170803
- Cicone F, Carideo L, Scaringi C et al (2019) F-DOPA uptake does not correlate with IDH mutation status and 1p/19q co-deletion in glioma. Ann Nucl Med 33(4):295–302. https://doi.org/10.1007/ s12149-018-01328-3
- Rao AM, Quddusi A, Shamim MS (2018) The significance of MGMT methylation in glioblastoma multiforme prognosis. J Pak Med Assoc 68(7):1137–1139

- Cimini A, Chiaravalloti A, Ricci M, Villani V, Vanni G, Schillaci O (2020) MGMT promoter methylation and IDH1 mutations do not affect [18F]FDOPA uptake in primary brain tumors. Int J Mol Sci. https://doi.org/10.3390/ijms21207598
- 73 Somme F, Bender L, Namer IJ, Noël G, Bund C (2020) Usefulness of 18F-FDOPA PET for the management of primary brain tumors: a systematic review of the literature. Cancer Imaging 20(1):70. https://doi.org/10.1186/s40644-020-00348-5
- 74 Tabatabaei P, Visse E, Bergström P, Brännström T, Siesjö P, Bergenheim AT (2017) Radiotherapy induces an immediate inflammatory reaction in malignant glioma: a clinical microdialysis study. J Neurooncol 131(1):83–92. https://doi.org/10.1007/ s11060-016-2271-1
- Tatekawa H, Uetani H, Hagiwara A et al (2021) Worse prognosis for IDH wild-type diffuse gliomas with larger residual biological tumor burden. Ann Nucl Med 35(9):1022–1029. https://doi.org/ 10.1007/s12149-021-01637-0
- 76. Janvier L, Olivier P, Blonski M et al (2015) Correlation of SUV-derived indices with tumoral aggressiveness of gliomas in static 18F-FDOPA PET: use in clinical practice. Clin Nucl Med 40(9):e429–e435. https://doi.org/10.1097/RLU.0000000000 000897
- 77. Bund C, Heimburger C, Imperiale A et al (2017) FDOPA PET-CT of nonenhancing brain tumors. Clin Nucl Med 42(4):250–257. https://doi.org/10.1097/RLU.000000000001540

- Patel CB, Fazzari E, Chakhoyan A et al (2018) F-FDOPA PET and MRI characteristics correlate with degree of malignancy and predict survival in treatment-naïve gliomas: a cross-sectional study. J Neurooncol 139(2):399–409. https://doi.org/10.1007/ s11060-018-2877-6
- 79 Mayerhoefer ME, Prosch H, Beer L et al (2020) PET/MRI versus PET/CT in oncology: a prospective single-center study of 330 examinations focusing on implications for patient management and cost considerations. Eur J Nucl Med Mol Imaging 47(1):51– 60. https://doi.org/10.1007/s00259-019-04452-y
- Ponisio MR, McConathy JE, Dahiya SM et al (2020) Dynamic. Neurooncol Pract 7(6):656–667. https://doi.org/10.1093/nop/ npaa044
- National Library of Medicine (U.S.) (2014) 18 F-DOPA- PET in Planning Surgery in Patients With Gliomas. Identifier NCT02020720. https://clinicaltrials.gov/ct2/show/NCT02020720
- Higgins GS (2021) [18F] FDOPA PET Imaging in Glioma: Feasibility Study for PET Guided Brain Biopsy. Identifier NCT04870580. https://clinicaltrials.gov/ct2/show/NCT04870580

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.