

Intracranial dissemination of glioblastoma multiforme: a case report and literature review

Keke Zhang¹, Yihang Yang², Jianfeng Zhuang³,
Gengyin Guo², Xin Chao⁴ and
Zhen Zhang^{2,5,6} 

Abstract

Intracranial dissemination is rare among patients with glioblastoma multiforme (GBM). Very few GBM patients develop symptoms from intracranial dissemination, as most do not surviving long enough for intracranial dissemination to become clinically evident. Herein, we report a case of GBM in a 39-year-old woman who underwent surgical resection, concomitant chemoradiotherapy, and seven courses of adjuvant chemotherapy with temozolomide. The patient then complained of an instable gait and hearing loss. Imaging studies demonstrated that although the primary intracranial tumors were well-controlled by treatment, contralateral cerebellopontine angle seeding dissemination was present. The patient died 3 months after the diagnosis of seeding dissemination. In light of a previous report and our current case, heightened awareness could promote surgical strategies that minimize the possibility of dissemination, including avoiding ventricular entry or a no-touch strategy.

Keywords

Glioblastoma, case report, intracranial dissemination, prognosis, no-touch strategy, temozolomide

Date received: 28 March 2022; accepted: 20 June 2022

¹Department of Otolaryngology, Shandong Provincial Hospital Affiliated to Shandong First Medical University

²Department of Neurosurgery, Shandong Provincial Hospital Affiliated to Shandong First Medical University
³Department of Neurosurgery, Qilu Hospital, Shandong University

⁴Department of Neurosurgery, Second People's Hospital of Xintai

⁵State Key Laboratory of Translational Medicine and Innovative Drug Development, Jiangsu Simcere Pharmaceutical Co., Ltd.

⁶The First Clinical College of Shandong University of Chinese Traditional Medicine

Corresponding author:

Zhen Zhang, Department of Neurosurgery, Shandong Provincial Hospital Affiliated to Shandong First Medical University, No. 324 Jingwuwei Road, Huaiyin District, Jinan City, Shandong Province, Jinan, 250021, China.
Email: zhangzhen@sdfmu.edu.cn



Introduction

Glioblastoma multiforme (GBM) is the most aggressive primary brain tumor in adults, accounting for 12% to 15% of all intracranial tumors.¹ The prognosis of GBM patients remains poor even with improvements in surgery, chemotherapy, and radiotherapy. During the last few years there have been an increasing number of reports about GBM dissemination, which is attributed to the prolonged survival of GBM patients after tumor excision.^{2,3} However, the incidence of GBM dissemination is lower than what is observed in autopsy studies.⁴ Many studies have shown that GBM dissemination induces fatal outcomes. The median survival time of GBM patients ranges from 11 to 17 months; however, the average time between a diagnosis of GBM seeding dissemination and death is only 2 to 3 months.⁵

Case report

A 39-year-old female patient with an unremarkable past medical history presented with headache and unilateral limb weakness and was admitted to our hospital. Neurological examination showed no positive signs. A head magnetic resonance imaging (MRI) scan revealed a long-T1 and long-T2 cystic and solid mass measuring $4.9 \times 4.7 \times 3.6$ cm with obvious peritumoral edema in the right frontotemporal lobe, which compressed the lateral ventricles. After gadolinium injection, the mass was heterogeneously enhanced (Figure 1a, b and c). Then the patient underwent a gross total resection of the right frontotemporal tumor. The methylation status of the O6-methylguanine-DNA methyltransferase (*MGMT*) promoter in the GBM sample was analyzed using methylation-specific PCR, and the *MGMT* promoter was found to be unmethylated.

Pathological examination of the tumor identified it as GBM. Immunohistochemistry (IHC) of the tumor demonstrated negative staining for glial fibrillary acidic protein (GFAP). The Ki-67 labeling index was 60% (Figure 2a, b, c and d). Three weeks after uneventful postoperative recovery, the patient started to receive concomitant chemoradiotherapy with temozolomide (TMZ; $75 \text{ mg/m}^2/\text{day}$) and external beam X-ray irradiation therapy (60 Gy in 30 fractions), which was tolerated well. Adjuvant chemotherapy with TMZ ($200 \text{ mg/m}^2/\text{day}$) for 5 days every 4 weeks was subsequently administered. The patient's initial symptoms were resolved, and head MRI 7 months after primary tumor resection showed no recurrence (Figure 1 d, f and g).

It should be noted that after the seventh cycle of adjuvant chemotherapy, the patient complained of an instable gait and hearing loss. Gadolinium-enhanced head MRI revealed a new heterogeneously-enhanced solid mass in the left cerebellopontine angle region but no recurrence of the initial tumor in the right frontotemporal lobe and no spinal dissemination (Figure 3e, f and g). The patient underwent secondary resection of the tumor, and histology confirmed GBM dissemination. IHC of the new mass demonstrated positive staining for GFAP and OLIG2 and a Ki-67 labeling index of 30% (Figure 3a, b, c, and d). Her family refused further treatment, and the patient died 3 months after the second surgery, 12 months after the initial diagnosis. We have also included a table summarizing previously published articles of related cases (Table 1). The reporting of this study conforms to CARE guidelines.⁶

Discussion

GBM is the most common malignant primary brain tumor, accounting for 12% to 15% of intracranial neoplasms.⁷ GBM has a reported global incidence of 2 to 3 per




Figure 1. Gadolinium-enhanced T1-weighted magnetic resonance imaging showing the intracranial tumors in the right frontal and insula prior to surgery (a: coronal image, b: sagittal image, c: axial image). Gadolinium-enhanced T1-weighted magnetic resonance imaging 7 months after surgery showing the tumor had been completely removed (d: coronal image, e: sagittal image, f: axial image).

100,000 people.¹ Despite advancements in diagnostic and therapeutic approaches, GBM continues to have a poor prognosis with patient survival at approximately 15 months after diagnosis.⁸ Previous studies indicated that GBM dissemination does not occur, primarily due to the strong protective mechanisms in the central nervous system, such as the lack of a true lymphatic system in the brain and because the venous sinuses are encased in dense dural membranes, both of which hinder invasion.^{9–11} GBM patients have a short survival time and commonly die from oncothlipsis, intracranial hypertension, and other complications before metastasis develops.¹²

Advancements in early detection and therapeutic approaches have resulted in an increased median survival for GBM patients, which has consequently increased

the detection of GBM dissemination.¹² Autopsy studies have also described that approximately 25% of GBM patients have evidence of spinal subarachnoid seeding, suggesting GBM dissemination is not uncommon.^{1,13,14} With the prolonged survival of GBM patients, it is important to study the GBM dissemination.

The most common metastasis sites for GBM are the spinal cord, lungs, bone, and lymph nodes.³ To date, the factors that cause GBM metastasis remain unclear. Cellular spread in the subarachnoid space seems to be the most likely cause for intracranial and spinal dissemination. According to the literature, ventricular entry at operation, repeated tumor resection, male sex, ependymal invasion, fissuring of the ependymal due to hydrocephalus, depressed immune function after radiotherapy and




Figure 2. (a) Hematoxylin and eosin staining of pathological sections after the first operation. (b) Immunohistochemistry showing the tumor was negative for GFAP; (c) the tumor was positive for OLIG2; (d) the Ki-67 labeling index was 60%.




Figure 3. (a) Hematoxylin and eosin staining of pathological sections after the second operation. (b) Immunohistochemistry showing the tumor was positive for GFAP; (c) the tumor was positive for OLIG2; (d) the Ki-67 labeling index was 30%. Magnetic resonance imaging performed after the 7th chemotherapy cycle revealed the presence of a mass in the left cerebellum. The mass appeared hyperintense on enhanced magnetic resonance imaging performed, with obvious heterogeneous enhancement (e: coronal image, f: sagittal image, g: axial image).

Table 1. Summary of previous cases reporting patients with GBM metastases.

Authors	Sex	Age at diagnosis	Tumor Location	Treatments after operation	Histopathological finding	Metastatic sites	Treatments after metastasis	Survival time after diagnosis of metastasis
L-T. Kuo et al. ¹⁶	M	41	left parieto-occipital lobe and another smaller lesion located anterior to the main mass	concurrent chemotherapy with TMZ and external beam X-ray irradiation therapy	glioblastoma multiforme	the entire spinal cord and brainstem	whole-spine RT	2 months
S. Battaglia et al. ¹⁷	M	11	intramedullary and exophytic mass extending from Th4 to Th5	RT to the whole spinal axis and TMZ	glioblastoma multiforme	spinal diffuse leptomeningeal and spinal root, a leptomeningeal brain and the left hippocampus	cerebrospinal fluid ventriculoperitoneal shunt	6 months
B.I. Ogunbo et al. ¹⁸	F	49	occipital lobe	chemotactic agents CCNU, procarbazine, metopoclopramide and RT	glioblastoma multiforme	left parotid gland	palliative oncological treatment	16 months
A. Mujic et al. ¹⁹	M	39	left frontal lobe	RT	glioblastoma multiforme	the left posterior parietal region, the pleura, small bowel, and pancreas	frame-less stereotactic excision	3 months
J.J. Grah et al. ²⁰	F	59	right frontal area	RT and concurrent chemotherapy	glioblastoma multiforme	the right frontal lobe, cervical leptomeningeal	second surgery, adjuvant RT and chemotherapy	1 months
I.J. Torres et al. ²¹	F	63	right upper frontal gyrus	cranial RT and TMZ	glioblastoma multiforme	the dura, scalp, and subcutaneous cell tissue	second surgery, carmustine	NA
S. Scocianti et al. ²²	M	33	right temporoparietal area	RT and TMZ, nitrosourea, and fotemustine	glioblastoma multiforme	intramedullary and leptomeningeal	palliative RT	4 months
T.K. Tshuhara et al. ²³	F	55	tip of middle fossa to the frontal base	RT and TMZ	glioblastoma multiforme	spinal cord	RT and chemotherapy	4 months
W. Zhang et al. ²⁴	M	47	left temporal lobe	RT and TMZ	glioblastoma multiforme	left frontotemporal lobe	N/A	16 months
G. Simonetti et al. ²⁵	M	38	left parietal lobe	RT and TMZ	glioblastoma multiforme	lung, lymph nodes, bones	local RT and chemotherapy	2 months
M. Taha et al. ²⁶	M	33	left frontal lobe	RT, chemotherapy, and resection of intracranial recurrence	glioblastoma multiforme	left parotid gland, cervical lymph nodes	local RT, PCV chemotherapy	3 months

Note: RT: radiation therapy; TMZ: temozolomide; CCNU: lomustine; PCV: procarbazine, lomustine, vincristine.

chemotherapy, and fragmentation of the tumor in contact with cerebrospinal fluid were all associated with a statistically significant increased incidence of central nervous system dissemination.³

In this case, the primary tumor was located at the lateral fissure, making cellular spread in the subarachnoid space the most likely cause. Heightened awareness can promote surgical strategies that minimize the possibility of dissemination, including avoiding ventricular entry or a no-touch strategy.

The prognosis of GBM patients with dissemination is bleak and almost always leads to fatal outcomes. The median time between a diagnosis of GBM dissemination and death is approximately 2 to 3 months,¹⁵ and treatment is primarily palliative.

Conclusion

GBM dissemination is common to some extent. Heightened awareness will promote surgical strategies that reduce the possibility of dissemination, including avoiding ventricular entry and a no-touch strategy.

Declaration of conflicting interest

The authors declare that there is no conflict of interest.

Ethics statement

Written informed consent was obtained from the parents for the publication of this case report. The study protocol was approved by Shandong Provincial Hospital affiliated to Shandong First Medical University.

Funding

The author(s) disclosed receipt of the following financial support for the research, authorship, and/or publication of this article: This work was supported by The Natural Science Foundation of Shandong Province (grant number: ZR2021MH228).

ORCID iD

Zhen Zhang  <https://orcid.org/0000-0002-6072-353X>

References

1. Stark AM, Nabavi A, Mehdorn HM, et al. Glioblastoma multiforme-report of 267 cases treated at a single institution. *Surgical neurology* 2005; 63: 162–169; discussion 169. 2005/02/01. DOI: 10.1016/j.surneu.2004.01.028.
2. Kong X, Wang Y, Liu S, et al. Brain Stem and Entire Spinal Leptomeningeal Dissemination of Supratentorial Glioblastoma Multiforme in a Patient during Postoperative Radiochemotherapy: Case Report and Review of the Literatures. *Medicine (Baltimore)* 2015; 94: e962. 2015/06/20. DOI: 10.1097/md.0000000000000962.
3. Lawton CD, Nagasawa DT, Yang I, et al. Leptomeningeal spinal metastases from glioblastoma multiforme: treatment and management of an uncommon manifestation of disease. *Journal of neurosurgery Spine* 2012; 17: 438–448. 2012/09/11. DOI: 10.3171/2012.7.Spine.12212.
4. Karaca M, Andrieu MN, Hicsonmez A, et al. Cases of glioblastoma multiforme metastasizing to spinal cord. *Neurology India* 2006; 54: 428–430. 2006/11/23. DOI: 10.4103/0028-3886.28122.
5. Grabb PA, Albright AL and Pang D. Dissemination of supratentorial malignant gliomas via the cerebrospinal fluid in children. *Neurosurgery* 1992; 30: 64–71. 1992/01/01. DOI: 10.1227/00006123-199201000-00012.
6. Gagnier JJ, Kienle G, Altman DG, et al. The CARE guidelines: consensus-based clinical case reporting guideline development. *Headache* 2013; 53: 1541–1547. 2013/11/26. DOI: 10.1111/head.12246.
7. Ray A, Manjila S, Hdeib AM, et al. Extracranial metastasis of glioblastoma: Three illustrative cases and current review of the molecular pathology and management strategies. *Molecular and clinical oncology* 2015; 3: 479–486. 2015/07/03. DOI: 10.3892/mco.2015.494.

8. Thakkar JP, Dolecek TA, Horbinski C, et al. Epidemiologic and molecular prognostic review of glioblastoma. *Cancer Epidemiol Biomarkers Prev* 2014; 23: 1985–1996. 2014/07/24. DOI: 10.1158/1055-9965.Epi-14-0275.
9. Figueroa P, Lupton JR, Remington T, et al. Cutaneous metastasis from an intracranial glioblastoma multiforme. *J Am Acad Dermatol* 2002; 46: 297–300. 2002/01/25. DOI: 10.1067/mj.2002.104966.
10. Al-Ali F, Hendon AJ, Liepman MK, et al. Oligodendroglioma metastatic to bone marrow. *AJNR Am J Neuroradiol* 2005; 26: 2410–2414. 2005/10/13.
11. Wu W, Zhong D, Zhao Z, et al. Postoperative extracranial metastasis from glioblastoma: a case report and review of the literature. *World J Surg Oncol* 2017; 15: 231. 2017/12/30. DOI: 10.1186/s12957-017-1300-7.
12. Lun M, Lok E, Gautam S, et al. The natural history of extracranial metastasis from glioblastoma multiforme. *J Neurooncol* 2011; 105: 261–273. 2011/04/23. DOI: 10.1007/s11060-011-0575-8.
13. Tantongtip D and Rukkul P. Symptomatic leptomeningeal and entirely intramedullary spinal cord metastasis from supratentorial glioblastoma: a case report. *J Med Assoc Thai* 2011; 94: S194–S197. 2012/05/25.
14. Raheja A, Borkar SA, Kumar R, et al. Metachronous spinal metastases from supratentorial anaplastic astrocytoma. *Asian J Neurosurg* 2015; 10: 60. 2015/03/15. DOI: 10.4103/1793-5482.151529.
15. Lomax AJ, Yannakou CK and Rosenthal MA. Spinal cord metastasis in a patient treated with bevacizumab for glioblastoma. *Target Oncol* 2013; 8: 153–155. 2013/02/23. DOI: 10.1007/s11523-013-0270-0.
16. Kuo LT, Tsai SY, Yang CY, et al. Meningeal seeding from glioblastoma multiforme treated with radiotherapy and temozolomide. *Asian J Surg* 2017; 40: 61–65. 2013/08/28. DOI: 10.1016/j.asjsur.2013.07.012.
17. Battaglia S, Albini Riccioli L, Bartiromo F, et al. Childhood spinal glioblastoma multiforme with intracranial dissemination. A case report. *Neuroradiol J* 2007; 20: 500–503. 2007/10/31. DOI: 10.1177/197140090702000504.
18. Ogungbo BI, Perry RH, Bozzino J, et al. Report of GBM metastasis to the parotid gland. *J Neurooncol* 2005; 74: 337–338. 2005/09/01. DOI: 10.1007/s11060-005-1480-9.
19. Mujic A, Hunn A, Taylor AB, et al. Extracranial metastases of a glioblastoma multiforme to the pleura, small bowel and pancreas. *J Clin Neurosci* 2006; 13: 677–681. 2006/07/04. DOI: 10.1016/j.jocn.2005.08.016.
20. Grah JJ, Katalinic D, Stern-Padovan R, et al. Leptomeningeal and intramedullary metastases of glioblastoma multiforme in a patient reoperated during adjuvant radiochemotherapy. *World J Surg Oncol* 2013; 11: 55. 2013/03/19. DOI: 10.1186/1477-7819-11-55.
21. Jusué Torres I, Jerez Fernández P, Ortega Zufiria J, et al. Skin spread from an intracranial glioblastoma: case report and review of the literature. *BMJ Case Rep* 2011; 2011: bcr0920114858. 2011/01/01. DOI: 10.1136/bcr.09.2011.4858.
22. Scoccianti S, Detti B, Meattini I, et al. Symptomatic leptomeningeal and intramedullary metastases from intracranial glioblastoma multiforme: a case report. *Tumori* 2008; 94: 877–881. 2009/03/10.
23. Katsuhara T, Moro N, Ohta T, et al. Solitary primary intracranial leptomeningeal glioblastoma invading the normal cortex: Case report. *Mol Clin Oncol* 2018; 8: 466–470. 2018/03/31. DOI: 10.3892/mco.2018.1561.
24. Zhang W, Cai YY, Wang XL, et al. Bone Metastases of Glioblastoma: A Case Report and Review of the Literature. *Front Oncol* 2021; 11: 705455. 2021/10/15. DOI: 10.3389/fonc.2021.705455.
25. Simonetti G, Silvani A, Fariselli L, et al. Extra central nervous system metastases from glioblastoma: a new possible trigger event? *Neurol Sci* 2017; 38: 1873–1875. 2017/06/26. DOI: 10.1007/s10072-017-3036-0.
26. Taha M, Ahmad A, Wharton S, et al. Extracranial metastasis of glioblastoma multiforme presenting as acute parotitis. *Br J Neurosurg* 2005; 19: 348–351. 2006/02/04. DOI: 10.1080/02688690500305506.