

Review

Implications of Advances in Studies of O⁶-Methylguanine-DNA-Methyltransferase for Tumor Prognosis and Treatment

Yuexia Chen¹, Wei Qu¹, Jianhong Tu¹, Hongyan Qi^{2,*}

¹Department of Pathology, The Third Hospital of Nanchang, 330008 Nanchang, Jiangxi, China

²Department of Pathology and Pathophysiology and Department of Radiation Oncology of the Second Affiliated Hospital, University School of Medicine, Zhejiang University, 310058 Hangzhou, Zhejiang, China

*Correspondence: qihongyan@zju.edu.cn (Hongyan Qi)

Academic Editor: Alexandros G. Georgakilas

Submitted: 3 November 2022 Revised: 11 January 2023 Accepted: 10 February 2023 Published: 12 September 2023

Abstract

O⁶-methylguanine-DNA-methyltransferase (MGMT) is a DNA repair enzyme, which reverses the alkylation of guanine O⁶ through direct transfer of the methyl group, maintains the gene stability and avoids tumor occurrence. Studies have shown that *MGMT* gene methylation, polymorphism and protein expression are involved in the process of various tumor development, such as colon cancer, gastric carcinoma, etc. *MGMT* gene promotes methylation, protein expression and enzyme activity from various tissues, which results in different effects on the prognosis of patients. MGMT promoter methylation is a positive factor for the prognosis of Glioblastoma (GBM), which can prolong overall survival and progression-free survival, reduce the resistance of tumor cells to temozolomide treatment, and improve the prognosis. The treatment of tumors based on MGMT focuses on three aspects: targeting MGMT to increase the sensitivity of alkylated drug therapy in tumors, immunotherapy combined with alkylated agents on tumor treatment, and treatment for patients with MGMT promoter non-methylation. Similarly, a number of studies have targeted MGMT to reduce alkylated agent resistance in other systems. Although numerous studies on MGMT in tumors have been reported, there are problems that need to be solved, such as selection and consensus of MGMT promoter methylation detection methods (CpG detection sites, cut-off value) and the treatment of MGMT non-methylated GBM patients, especially elderly patients. In this review, we describe the regulation of MGMT expression and its role in chemotherapy, especially in gliomas. Further studies exploring new methods targeting MGMT with better curative effect and less toxicity are advocated. We anticipate that these developments will be progressive and sufficiently used for clinical application.

Keywords: MGMT; expression; prognosis; treatment; tumor

1. Introduction

O⁶-methylguanine-DNA-methyltransferase (MGMT), located on chromosome 10q26, is over 170 kb in length, including 5 exons and 4 introns. mRNA is 866 bp in length, encodes 207 amino acids and contains a conserved active site [1]. The promoter region is approximately 1.2 kb and contains enhancers, binding sites for transcription factors such as GR, Sp1, AP1, AP2 [2–4]. The promoter region lacks TATA and CAAT boxes, but is rich in GC sequences, forming 98 CpG sites. These sites are where cytosines are easily methylated, especially in the regions –186 to –172 and +93 to +153 where CpG methylation plays a major role in transcriptional regulation. MGMT expression is mainly regulated by transcriptional and epigenetic regulation, the former including SP1, AP1, NF-κB, GRE, p53, etc., while the latter contains DNA methylation in the promoter region and post-translational modifications of histones [5]. With cytosine methylation, the chromosome structure changes from loose and active euchromatin to compressed and aggregated heterochromatin, which prevents transcription factors from binding to the promoter region and inhibits transcription. The amino termini of histones could be modified by methylation,

acetylation, ubiquitination, and poly-ADP-ribosylation, and modification of H3K9 was associated with MGMT silencing [6]. With DNA methylation, methylated CpG island binding proteins, especially MeCP2, bind to the methylated CpG island and then recruit histone deacetylases and H3K4 demethylases (such as LSD1), as well as histone methyltransferase to form heterochromatin like protein factors (HP1), eventually preventing the transcription process [5]. miRNAs, such as miR-181b, miR-181d, miR-221, miR-222, miR-767-3p and miR-648n [7–10], bind to the 3' untranslated region of MGMT to reduce mRNA stability and affect protein translation.

MGMT is a highly conserved enzyme involved in DNA damage repair with a conserved amino acid sequence-(I/V) PCHR (Proline, cysteine, histidine, arginine) (V/I) located at the active center. During repair, MGMT independently transfers methyl groups from the guanine O⁶ site directly to its own 145 cysteine residue, which is then accompanied by irreversible ubiquitination degradation [11] (Fig. 1). If O⁶MeG is not repaired, it leads to a G:C → A:T mutation that can be recognized by the mismatch repair system (MMR), which then initiates an apoptotic signaling pathway, leading to an ineffective replication cycle, DNA



fragmentation, and eventually apoptosis [12]. MGMT can repair various types of damage caused by alkylated agents from a wide range of sources, including endogenous and exogenous damage, the latter of which encompasses the DNA causing methylation damage at the guanine O⁶ site, such as temozolomide (TMZ), and drugs that cause O⁶-CLG damage at the guanine O⁶ site on DNA, such as carmustine mustard (BCNU, BiCNU), lomustine (CCNU and CeeNU) and other chemotherapeutic agents [13,14]. MGMT transfers methyl from the guanine O⁶ site directly to its cysteine residue at position 145 to complete repairing O⁶-mG damage of DNA.




Fig. 1. The process of (MGMT) repairing damaged DNA. MGMT, O⁶-methylguanine-DNA-methyltransferase; Cys, Cysteine.

2. Role of MGMT in Tumorigenesis and Progression

The G:C → A:T mutation in *K-ras*, *P53*, and other oncogenes or tumor suppressor genes is caused by MGMT down-regulation of protein expression as a result of promoter methylation, which favors the emergence and growth of malignancies. With this modification, a clear connection has been made between MGMT decrease and tumorigenesis [15]. In gastric cancer, colorectal cancer, breast cancer, oral squamous cell carcinoma and cervical squamous cell carcinoma tissues, MGMT promoter methylation is higher than that of surrounding normal mucosa or tissues. Therefore, MGMT promoter methylation might be considered a marker of precancerous lesions and a biomarker for early tumor diagnosis [16–19]. Another meta-analysis revealed that MGMT promoter hypermethylation was strongly linked to a higher risk of developing gastric cancer and may be connected to the spread of gastric cancer to distant sites and lymph nodes [20]. In our previous study to explore the molecular mechanism of MGMT in the malignant transformation and tumorigene-

sis of gastric cells induced by amide compounds, we found that MGMT up-regulation was induced by promoter hypomethylation. High expression of MGMT can prevent malignant transformation and tumorigenesis induced by amide compounds. In addition, in normal gastric tissues and gastric cancer patient specimens, MGMT was up-regulated in precancerous lesions and metaplastic tissues, and down-regulated in gastric cancer tissues, suggesting that MGMT may be involved in the occurrence and development of gastric cancer [21]. Additionally, MGMT has been shown to have a role in the development of other pathologies such as liver cancer, cholangiocarcinoma, lung cancer and other tumors [22–24]. *MGMT* gene polymorphisms have also been shown to be involved in tumor formation. A Mexican study demonstrated that MGMT rs12917 may contribute to the occurrence and progression of lung cancer [25]. It has also been revealed that women having the *MGMT* gene polymorphism Ile143Val have a lower chance of developing rectal cancer [26]. Single nucleotide polymorphisms in the *MGMT* gene can also affect the duration of TMZ-induced myelotoxicity and the side effects of antitumor drugs in adult patients with diffuse glioma [27]. MGMT polymorphism rs12917 might affect the response to chemotherapy in pediatric Hodgkin lymphoma [28]. Several investigations have discovered a link between the MGMT V1/W genotype and glioma recurrence [29] with other studies on the association between *MGMT* gene polymorphisms and tumors being performed.

3. Effect of MGMT on Tumor Prognosis

3.1 Effect of MGMT Promoter Methylation on Prognosis of Different Tumors

MGMT promoter methylation varied among different tumors. Approximately 38% of brain tumors were prone to MGMT promoter methylation, as were 28% of head and neck tumors, 26% of colon cancer, 25% of lymphoma, and 24% of lung cancer, while tumors of other organs exhibited a lower incidence of MGMT promoter methylation, such as pancreatic cancer, melanoma, kidney cancer, bladder cancer, and leukemia [30].

Numerous studies have demonstrated that MGMT promoter methylation is a beneficial factor for tumor prognosis. It has been proven to be related to improved patient prognosis, particularly in Glioblastoma (GBM), as well as prolonged overall survival (OS) and progression-free survival (PFS).

In a prospective clinical experiment conducted in 2004, Hegi *et al.* [31] showed the prognostic significance of MGMT promoter methylation for the prognosis of GBM patients receiving TMZ, an alkylating drug. The study revealed that MGMT promoter methylation leading to *MGMT* gene inactivation was associated with longer survival in GBM patients, suggesting that MGMT promoter methylation remains the only significant predictor for GBM patients [31]. In a subsequent study of 206 GBM patients, the re-

searchers found that GBM patients with MGMT promoter methylation responded to TMZ, while those who did not were more likely to develop treatment resistance [32]. Subsequently, other researchers have carried out a large number of studies on the correlation between MGMT promoter methylation and prognosis of GBM patients and found the prognosis of grade II and III gliomas to be positively impacted by MGMT promoter methylation [33,34].

Szylberg *et al.* [35] collected data on 41 newly diagnosed GBM patients treated at the 10th Military Research Hospital and Polyclinic in Poland between 2011 and 2014 to assess the impact of MGMT promoter methylation and other clinical factors on survival in patients with primary GBM (mean age, 53 years). They discovered that MGMT promoter methylation, which may be correlated with age and surgical resection technique, was also a significant positive predictive biomarker in GBM patients [35].

The results are still debatable even though the prognostic value of MGMT promoter methylation in GBM patients are well acknowledged. Some studies have not shown a significant correlation between MGMT promoter methylation and OS or PFS. In a multicenter Portuguese investigation, neither univariate nor multivariate analysis revealed a statistically significant relationship between MGMT promoter methylation and overall survival or PFS in patients with GBM receiving TMZ-based chemoradiotherapy [36]. Egaña *et al.* [37] also concluded that MGMT promoter methylation did not affect patient survival in the cohort studied. However, around 9 months after the diagnosis of GBM, Dahlrot *et al.* [38] discovered the connection between MGMT promoter methylation and overall survival, as prior to this, there was no correlation between MGMT promoter methylation and OS. In addition, Caccese *et al.* [39] found there was a nonlinear correlation between MGMT promoter methylation and OS, with longer OS with increasing MGMT promoter methylation. This study showed a median OS of 14.8 months for MGMT 0–4%, 18.9 months for MGMT 4–40%, and 29.9 months for MGMT 40–100% [39]. MGMT promoter methylation has also been used to stratify malignant gliomas in GBM and has been used as a prognostic marker for overall survival and as a predictor of chemotherapy response in GBM patients [40].

Numerous studies have been conducted in other tumors in addition to the one on MGMT promoter methylation and GBM, and the effects of tumor prognosis varied. MGMT promoter methylation in succinate dehydrogenase defect of gastrointestinal stromal tumors and epithelioid/mixed phenotype is particularly common in the wild type of gastrointestinal stromal tumor and this MGMT methylation pattern may provide a new potential treatment option for wild-type gastrointestinal stromal tumor [41]. In well-differentiated pancreatic neuroendocrine tumors, PFS is longer in patients with high MGMT promoter methylation and low MGMT expression [42]. In metastatic colorec-

tal cancer, melanoma, central nervous system lymphoma and other tumors, MGMT promoter hypermethylation increased the sensitivity of alkylating agents, thus prolonging the survival of patients [43–45]. Numerous studies in esophageal, cervical, and lung cancers have shown controversial results [24,46,47]. These differences may be caused by: differences in MGMT promoter methylation detection assays or different methylation sites in the detected MGMT promoter region, different samples, different numbers contained in different tumor cells, and patient age [48].

MGMT gene polymorphism was also involved in predicting the prognosis of many tumors. Researchers have found that *MTNR1B* single-nucleotide polymorphisms (SNPs) combined with *CDKN2A* and MGMT promoter methylation status can be used to predict shorter survival in colon cancer [49].

3.2 Effect of MGMT Protein Expression on Prognosis of Different Tumors

MGMT, a DNA repair enzyme, is expressed in many organs of the body and its expression varies from one organ and tissue to another. The expression of *MGMT* protein was highest in the liver, followed by lung, kidney and colon, and lowest in the pancreas, hematopoietic cells, lymphoid tissues and brain. It was decreased in tumors such as gliomas, lymphomas, breast cancer, prostate cancer, and retinoblastoma, most likely related to the methylation status of its promoter region [50].

While *MGMT* protein expression detection methods mainly include immunofluorescence and immunohistochemistry (IHC), the latter predominates. In the early 1990s, Belanich *et al.* [51] used immunofluorescence method to detect the expression of *MGMT* protein in 99 cases of glioma tissues and found that patients with high *MGMT* protein expression were less sensitive to BCNU, resulting in shorter overall survival and progression-free survival. This was subsequently demonstrated in astroglomas [51]. With various tumor types, there are differences in the expression of the *MGMT* protein as identified by the IHC approach and the prognosis of the tumor. A report of 73 cases of patients with newly diagnosed GBM utilized immunohistochemical methods to analyze *MGMT* protein expression as an auxiliary for TMZ and radiation treatment of GBM. In patients with detected prognostic markers, it was found that low *MGMT* protein expression in patients (15%) compared with patients with high expression of *MGMT* protein, overall survival and PFS were significantly improved. This suggests that *MGMT* protein expression is an independent and a positive prognostic factor in GBM patients [52]. *MGMT* immunohistochemistry expression has been demonstrated to be substantially correlated with various glioma grades and subtypes [53] including lymphomas, thymic tumors, and pituitary tumors [54–56].

Other investigations have reached the opposite conclusion. Aanchal *et al.* [57] collected the tissues of pa-

tients with meningeal hemangiopericytoma from 2002 to 2011. Immunohistochemistry was used to detect expression of MGMT protein and methylation-specific PCR (MSP) was adopted to detect MGMT promoter methylation. In investigating the relationship between MGMT expression and the prognosis of patients with meningeal hemangiopericytoma, they found no significant correlation between MGMT protein expression and progression-free survival [57]. In esophageal and gastrointestinal neuroendocrine tumors, MGMT protein expression level was also shown to have no specific correlation with prognosis, whereas in pancreatic neuroendocrine tumors, colorectal cancer and salivary gland carcinoma, patients with low MGMT protein expression had a poorer prognosis [58–61]. A recent study found that decreased MGMT expression in pancreatic neuroendocrine tumors was associated with a higher risk of progression [62]. The reasons for these discrepancies may be related to tumor type, the number of patients studied, individual differences, tissue size, medication differences, and diagnostic staining.

3.3 Effect of MGMT Enzyme Activity on Prognosis of Different Tumors

The enzyme activity of MGMT varies with different tissues, individuals and individual stages. In normal tissues, MGMT enzyme activity in the liver was highest, and lowest in brain tissue. While MGMT activity was highest in liver, ovarian, and colon tumors, it was still very low in gliomas, which may have contributed to the sensitivity of glioma cells to TMZ therapy. The activity varied according to the classification of gliomas, being lowest in astrocytomas and malignant gliomas, with an average of 111 fmol/mg, and up to 270 fmol/mg in non-glioma brain tumors, such as meningiomas [63]. MGMT activity in tumor tissues was higher than that in corresponding normal organ tissues, but in the testes and liver, the activity of tumor tissues was lower than that in normal tissues. There is minimal research examining the connection between MGMT activity and tumor prognosis. Early research has suggested that elevated MGMT activity may reduce the effectiveness of alkylated anticancer medications and have an adverse effect on patient prognosis [64,65].

4. Effect of MGMT on Tumor Therapy

MGMT is a DNA repair that inhibits the cross-linking of double-stranded DNA through alkylated agents, reverses the alkylation of guanine O⁶ position, repairs DNA damage caused by drugs (such as alkylating agents), and leads to resistance to alkylated drugs [66]. It is also involved in the resistance of DNA to alkylated anticancer drugs such as TMZ, which has been studied extensively to overcome these therapeutic difficulties.

4.1 Targeting MGMT to Increase the Sensitivity of TMZ Therapy in Tumors

Although previous clinical trials have enhanced the therapeutic effect of TMZ by reducing MGMT protein expression, certain trials did not achieve clinical benefits [67]. A growing number of studies have focused on increasing the sensitivity of tumors (especially GBM) to TMZ treatment by targeting MGMT through various axes. Geng *et al.* [68] found that exosome-mediated circWDR62 promoted TMZ resistance and progression in glioma by targeting the Mir-370-3p/MGMT axis, suggesting that exosomal circWDR62 in human serum may be a therapeutic target for glioma. Zhou *et al.* [69] found that cyanidin-3-o-glucoside promoted the treatment of MGMT-induced glioma cell resistance through the potential signaling mechanism of miR-214-5p-mediated inhibition of TMZ resistance in LN-18/TR cells. Other studies have demonstrated that the lncRNA UCA1/miR-182-5p/MGMT axis regulates the sensitivity of glioma cells to TMZ through the MGMT-related DNA damage pathway [69,70]. In addition to gliomas, other tumors that have been studied include melanoma, lymphoma, and ovarian cancer [71–73].

Other potential targets have been investigated. The MGMT substrate analogue, O⁶-benzylguanine, was a specific inhibitor of MGMT, which binds the benzoyl group to the 145th cysteine residue in the active center of MGMT protein, preventing the binding of the latter to DNA. It effectively reduces the ability of MGMT to repair alkyl-adduct DNA and inhibits MGMT activity. However, it has never been used clinically due to its side effects [74]. Pinto *et al.* [75] designed a hybrid drug that covalently binds BG residues to the interacting part of DNA (6-chloro-2-methoxy-9-aminoacridine) and found that compound 19a inhibits MGMT activity without inducing significant levels of DNA damage, providing a new therapeutic opportunity for GBM patients with TMZ resistance.

Numerous studies have investigated MGMT and tumor medication resistance by manipulating additional upstream and downstream signaling pathways in addition to the use of the aforementioned inhibitors. A study from China found that BanxiaXiexin decoction regulates MGMT expression through IL6/JAK/STAT3-mediated PDL1 activity, which affects the sensitivity of gastric cancer cells to drugs. As a result, inhibiting MGMT offers a fresh approach to treating gastric cancer [76]. Additionally, it has been demonstrated that MGMT has a role in the chemosensitivity of cisplatin in gastric cancer [77]. MAPK/ERK inhibitor (U0126) combined with TMZ can be used in patients with advanced hepatocellular carcinoma as the former can block the MAPK/ERK signaling pathway to down-regulate MGMT expression and increase the sensitivity of HCC cells to TMZ [78]. MGMT was involved in the resistance process of dacarbazine treatment in uveal melanoma, but TRIM72 increased the sensitivity of dacarbazine treatment by ubiquitination and degradation of

Table 1. A collection of anti-tumor agents targeting MGMT.

Compound	Mechanism	Tumor type	Ref
circWDR62	Targeting miR-370-3p/MGMT axis promotes TMZ resistance	glioma	Geng <i>et al.</i> [68]
Cyanidin-3-O-glucoside	Down-regulation of β -catenin and MGMT by miR-214-5p inhibited TMZ resistance	glioma cell	Zhou <i>et al.</i> [69]
lncRNA UCA1	lncRNA UCA1/miR-182-5p/MGMT axis modulates glioma cell sensitivity to TMZ	glioma cell	Cheng <i>et al.</i> [70]
lncRNA POU3F3	lncRNA POU3F3/miR-650 axis upregulates MGMT expression to promote drug resistance	melanoma	Wu <i>et al.</i> [71]
miR-370	miR-370 inhibited MGMT expression and increased sensitivity to TMZ	primary central nervous system lymphoma	Li <i>et al.</i> [72]
MCL1	HDACis combined with PaTrin-2 overcomes resistance via the MGMT-DUB3-MCL1 axis	ovarian cancer	Wu <i>et al.</i> [73]
NCT503	Modulation of Wnt/ β -catenin axis reduces MGMT expression to overcome drug resistance	glioblastoma	Jin <i>et al.</i> [80]
Tubeimoside-I (TBMSI)	Induced apoptosis in GBM cells through reducing MGMT expression and inhibiting the EGFR induced PI3K/Akt/mTOR/NF- κ B signaling pathway	glioblastoma multiforme	Tang <i>et al.</i> [81]
GNA13	GNA13/PRKACA/MGMT modulates glioma sensitivity to TMZ	glioma	Liu <i>et al.</i> [82]
Pyrrinium pamoate	The AKT/GSK3 β / β -catenin signaling axis regulates MGMT expression	glioblastoma	Li <i>et al.</i> [83]
DEC1	Control TMZ resistance via the SP1-MGMT axis	glioma cell lines	Lv <i>et al.</i> [85]
METTL3	Silencing MeTTL3-mediated inhibition of total methylation improves TMZ resistance	glioblastoma cell lines	Shi <i>et al.</i> [84]
Banxia xiexin decoction	Influence the drug sensitivity of GC cells by regulating the expression of MGMT	gastric cancer cells	Feng <i>et al.</i> [76]
Cisplatin (DDP)	Inhibits MGMT-mediated autophagy suppression to decrease chemosensitivity in GC	gastric cancer	Lei <i>et al.</i> [77]
U0126	Downregulate MGMT expression via blocking MAPK/ERK pathway increased sensitivity to TMZ	hepatocellular carcinoma cells	Li <i>et al.</i> [78]
Acridine-O ⁶ -benzylguanine hybrids	Compound 19a inactivates MGMT to increase TMZ sensitivity	glioblastoma cell lines	Franco <i>et al.</i> [75]

TMZ, temozolomide; GBM, Glioblastoma; GC, guanine-cytosine.

MGMT [79]. NCT503, Tubeimoside-I, GNA13, Pyrrinium pamoate, DEC1, METTL3, MMR have been confirmed to increase the sensitivity of GBM to TMZ treatment by regulating MGMT [80–86], as summarized in (Table 1, Ref. [68–73,75–78,80–85]).

4.2 Immunotherapy Combined with Alkylating Agents for Tumor Treatment

In recent years, immunotherapy has gained popularity as a cancer treatment option, although certain clinical trials have not produced positive outcomes. In phase 3 trials of TMZ plus nivolumab in newly diagnosed MGMT promoter methylated GBM, radiation therapy (RT) + TMZ plus the immune checkpoint inhibitor nivolumab did not improve survival [87]. Absent longer OS was seen compared to RT

plus TMZ in a further phase 3 study of TMZ with nivolumab in GBM patients lacking MGMT methylation [88].

4.3 Treatment of MGMT Promoter Non-Methylated GBM

Different randomized trials have shown that GBM patients with MGMT promoter methylation were associated with significantly higher survival when treated with combined radiotherapy and TMZ. Even when disease relapses, TMZ was increasingly beneficial for GBM patients with MGMT promoter methylation [89]. However, TMZ therapy was ineffective and had a dismal prognosis for MGMT non-methylated individuals.

Barazzuol *et al.* [90] found at the cytological level that PARP2, an important enzyme involved in DNA repair, and its inhibitor ABT-888, when used in combination with

X-ray and TMZ, could enhance radio sensitivity and chemical sensitivity in MGMT methylated cell lines as well as in MGMT non-methylated cell lines. These findings suggest that ABT-888, in combination with traditional chemotherapy and radiation, has the clinical potential to enhance the currently recommended course of treatment for GBM, particularly for individuals who lack MGMT methylation [90]. Jue *et al.* [91] tested the PARP inhibition of veliparib (ABT-888) in a group of patient-derived cell lines and patient-derived xenograft models, which confirmed that the use of veliparib and RT was an effective treatment for GBM patients with non-methylated MGMT promoter. In a randomized phase II trial involving patients with newly diagnosed MGMT-unmethylated GBM, Veliparib treatment was generally well tolerated, although there was insufficient proof for a therapeutic benefit [92]. In addition to the use of PARP inhibitors, Meclofenamate, a nonsteroidal anti-inflammatory drug, as a gap junction inhibitor, could enhance the vulnerability of GBM cells to lomustine induced cell death. However, it did not depend on the methylation status of the MGMT promoter, which provides a new therapeutic possibility for GBM patients without MGMT promoter methylation [93]. Also, miRNA has been shown to increase the sensitivity of MGMT promoter non-methylated GBM patients to TMZ. Kirstein *et al.* [94] found that miRNAs may be a promising and innovative treatment to improve TMZ sensitivity and increase progression-free and long-term survival in MGMT non-methylated GBM patients.

The survival benefit of a 70% degree of resection threshold in GBM patients with non-methylated MGMT promoter supports the maximization of safe resection rather than the “all or nothing” approach, according to research by Katsigiannis *et al.* [95]. Based on a study of 175 newly diagnosed patients with primary GBM [96], MGMT promoter methylation combined with complete resection was an independent predictor for improved overall and PFS in newly diagnosed isocitrate dehydrogenase (IDH) wild-type GBM. These studies suggest that the effect of the scope of surgical resection on the prognosis of GBM patients may be related to the methylation level of the MGMT promoter.

The most widely used prognostic procedures for GBM exclude individuals older than 70 as it is perceived that older patients have a decreased tolerance to surgery and chemotherapeutic treatments and have a poorer prognosis [97,98]. When older GBM patients over 70 years old were combined with MGMT promoter non-methylation, the treatment was more difficult. Yuen *et al.* [99] analyzed the historical evidence-based data of GBM treatment in older patients and proposed that for these patients with MGMT promoter methylation, single-dose TMZ may be considered, while for older patients with MGMT promoter non-methylation, subfractionated radiotherapy alone may be sufficient (Table 2, Ref. [90,91,93–95,99,100]).

Moreover, MGMT expression may affect the chemotherapy drugs and lessen their therapeutic effect. At the same time, chemotherapy drugs also affect MGMT expression. Numerous investigations have revealed that MGMT methylation or activity in patients with GBM altered after receiving chemotherapy and that the expression of MGMT in certain patients with recurrence was different from that of the initial tumor. No mechanisms have been elucidated in these phenomena or in correlating with the selectivity of chemotherapeutic agents to cells with high MGMT expression [101].

5. New Progress in Other Aspects of MGMT

There have been many other aspects of research on MGMT, such as the effect of MGMT promoter methylation status combined with other genes on tumor prognosis or treatment, along with the update of MGMT methylation detection methods.

In addition to the single factor of MGMT promoter methylation status affecting tumor prognosis, the combination with other genes can also affect tumor prognosis, especially for GBM patients. According to certain studies, GBM patients' tumors had OS and PFS that were considerably greater than those of patients with wild-type *IDH1* GBM tumors that had an unmethylated MGMT promoter [102,103]. In addition to *IDH1* mutation, some researchers have studied the effect of *TERT* promoter mutation status and MGMT promoter methylation status of different patterns on the prognosis of GBM, but the results have been conflicting. Some studies have suggested that *TERT* promoter mutation combined with MGMT promoter methylation can prolong overall survival and progression free survival [104,105]. The prognosis of GBM with MGMT promoter methylation and *TERT* wild-type was found to be superior to that of other subtypes in another study [106]. This discrepancy might be brought on by various MGMT promoter methylation detection techniques employed in various laboratories, individual variations, or sample size.

Methylation-specific PCR, pyrosequencing, or methylation arrays are recommended for detecting MGMT promoter methylation assays according to the European Association of Neuro-Oncology guidelines while the European Society for Medical Oncology does not recommend immunohistochemistry to determine MGMT promoter methylation status [107]. Earlier studies have shown that IHC assays are not recommended for evaluating MGMT status [108]. Subsequently, pyrosequencing has also been proven to be an ideal choice for detecting MGMT promoter methylation status, with a recommended biological cutoff of 10% or 21% of the receiver operating characteristic [109–111]. In addition, it has recently been found that pyrosequencing (PSQ) to evaluate the percentage of MGMT promoter methylation is very important for predicting the volume response and prognosis of patients with residual tumor GBM [112]. Other methods used to detect MGMT promoter methylation include Lab-on-Chip com-

Table 2. MGMT non-methylated GBM treatment.

Treatment	Mechanism	Tumor type	Ref
Veliparib (ABT-888)	induced apoptosis and decreased cell proliferation in a PDX of MGMT unmethylated GBM	glioblastoma	Jue <i>et al.</i> [91]
ABT-888	might be significant in MGMT-unmethylated patients less benefit from TMZ	glioblastoma	Barazzuol <i>et al.</i> [90]
Meclofenamate	reinforces the antitumoral effects of chemotherapeutic agent lomustine, independent of MGMT promoter methylation status	glioblastoma	Schneider <i>et al.</i> [93]
miRNAs	enhance TMZ sensitivity in MGMT unmethylated patients	glioblastoma	Kirstein <i>et al.</i> [94]
70% extent of resection threshold	benefit for survival of patients with MGMT unmethylated GBM	glioblastoma	Katsigiannis <i>et al.</i> [95]
Chemotherapy drugs	combination of carelizumab, anlotinib, and oxitinib	glioblastoma	Wang <i>et al.</i> [100]
hypofractionated radiation therapy (hRT)	hRT alone can be considered	elderly patients with newly diagnosed glioblastoma	Yuen <i>et al.</i> [99]

PDX, patient-derived xenograft.

patible isothermal amplification, two-probe quantification of MSP, and methylation quantification of endonuclease resistant DNA [113–115]. IHC assays for MGMT protein or enzyme activity have also been used as a proxy for methylation status, especially in resource-limited settings, where PCR technology is lacking [116]. The MGMT promoter methylation, methylation site preference, and cut-off value could not be determined using a conventional approach, regardless of the detection method used.

The identification of methylation locations and threshold values continue to be established, despite evidence linking MGMT promoter methylation status to therapy and prognosis in GBM patients. Research has found that a cut-off of 9% for 74–78 CpG sites is better than a higher cutoff of 28% or 29% [110]. As the MGMT promoter methylation detection PSQ threshold depends on the average methylation CpGs threshold, the recommended value was 10%, which was divided into “transition zone” or “gray area”, as it might confer some sensitivity to TMZ treatment [117]. IHC has proved to be a robust method for predicting the prognosis of patients in the gray area defined by PSQ [118].

Various researchers have turned their attention to radiomics imaging methods based on nuclear magnetic resonance imaging, trying to establish a preoperative, non-invasive MGMT promoter methylation detection [119]. Extreme Gradient Boosting feature selection model, genetic algorithm based packaging model, machine learning, intravoxel incoherent motion (IVIM) and dynamic susceptibility contrast (DSC), and T2-weighted image have all been proven to be effective in evaluating MGMT promoter methylation in GBM patients [120–124]. Patients can combine imaging prediction with surgical specimen detection. This combination can more precisely predict MGMT methylation levels and let patients choose a more

effective treatment strategy. There have been other studies that used patient body fluids to detect MGMT promoters’ methylation status, such as peripheral blood and cerebrospinal fluid [125,126].

6. Conclusions

In its capacity as a DNA repair enzyme, MGMT is independently involved in DNA damage repair and is crucial for the emergence and growth of malignancies. In the early stage of tumor development, the transfer methyl group of MGMT avoids gene mutations of oncogenes and tumor suppressor genes such as *K-ras*, *P53* and *PTEN*, and plays an anti-tumor role. MGMT performs a function in transferring the methyl group to protect the body throughout tumor formation when chemotherapy medications, primarily alkylated chemotherapeutic agents, are used. While MGMT increases the protective effect on cells, it also produces drug resistance, which may affect the patient’s prognosis.

Although there are numerous studies on the influence of MGMT promoter methylation, protein expression, enzyme activity and gene polymorphism on tumor prognosis and treatment, the results have been controversial. Problems to be solved, especially for GBM patients, include the lack of consensus on the selection of MGMT methylation detection methods, the selection of methylation sites, and the determination of cut-off values. The use of different methods and standards in various laboratories can lead to inconsistent results. In addition, TMZ resistance caused by MGMT is still the key to the treatment of GBM, and the development of more targeted drugs that could be combined with TMZ or inhibit MGMT expression is also critical, especially for older patients who are more than 70 years old, as within this vulnerable group, it is urgent to develop more effective treatment regimens.

Abbreviations

MGMT, O⁶-methylguanine-DNA-methyltransferase; GBM, Glioblastoma; OS, Overall survival; PFS, Progression free survival; TMZ, temozolomide; IHC, Immunohistochemistry.

Author Contributions

YC and HQ had the idea for the article; WQ and JT performed the literature search and drafted the work; and HQ critically revised the work. All authors reviewed the manuscript. All author read and approved the final manuscript.

Ethics Approval and Consent to Participate

Not applicable.

Acknowledgment

Not applicable.

Funding

This research was supported by Zhejiang Medical and Health Science and Technology Project (2022KY024), Zhejiang Provincial Natural Science Foundation of China (LY18H160024 and LY20H160040).

Conflict of Interest

The authors declare no conflict of interest.

References

- [1] Nakatsu Y, Hattori K, Hayakawa H, Shimizu K, Sekiguchi M. Organization and expression of the human gene for O⁶-methylguanine-DNA methyltransferase. *Mutation Research*. 1993; 293: 119–132.
- [2] Biswas T, Ramana CV, Srinivasan G, Boldogh I, Hazra TK, Chen Z, *et al.* Activation of human O⁶-methylguanine-DNA methyltransferase gene by glucocorticoid hormone. *Oncogene*. 1999; 18: 525–532.
- [3] Harris CC. Molecular basis of multistage carcinogenesis. *Princess Takamatsu Symposia*. 1991; 22: 3–19.
- [4] Boldogh I, Ramana CV, Chen Z, Biswas T, Hazra TK, Grösch S, *et al.* Regulation of expression of the DNA repair gene O⁶-methylguanine-DNA methyltransferase via protein kinase C-mediated signaling. *Cancer Research*. 1998; 58: 3950–3956.
- [5] Soejima H, Zhao W, Mukai T. Epigenetic silencing of the MGMT gene in cancer. *Biochemistry and Cell Biology*. 2005; 83: 429–437.
- [6] Dillon N. Heterochromatin structure and function. *Biology of the Cell*. 2004; 96: 631–637.
- [7] Wang J, Sai K, Chen FR, Chen ZP. miR-181b modulates glioma cell sensitivity to temozolomide by targeting MEK1. *Cancer Chemotherapy and Pharmacology*. 2013; 72: 147–158.
- [8] Zhang W, Zhang J, Hoadley K, Kushwaha D, Ramakrishnan V, Li S, *et al.* miR-181d: a predictive glioblastoma biomarker that downregulates MGMT expression. *Neuro-oncology*. 2012; 14: 712–719.
- [9] Quintavalle C, Mangani D, Roscigno G, Romano G, Diaz-Lagares A, Iaboni M, *et al.* MiR-221/222 target the DNA methyltransferase MGMT in glioma cells. *PLoS ONE*. 2013; 8: e74466.
- [10] Kreth S, Limbeck E, Hinske LC, Schütz SV, Thon N, Hoefig K, *et al.* In human glioblastomas transcript elongation by alternative polyadenylation and miRNA targeting is a potent mechanism of MGMT silencing. *Acta Neuropathologica*. 2013; 125: 671–681.
- [11] Pegg AE. Repair of O(6)-alkylguanine by alkyltransferases. *Mutation Research*. 2000; 462: 83–100.
- [12] Wick W, Weller M, van den Bent M, Sanson M, Weiler M, von Deimling A, *et al.* MGMT testing—the challenges for biomarker-based glioma treatment. *Nature Reviews. Neurology*. 2014; 10: 372–385.
- [13] Herman JG, Baylin SB. Gene silencing in cancer in association with promoter hypermethylation. *The New England Journal of Medicine*. 2003; 349: 2042–2054.
- [14] Christmann M, Kaina B. O(6)-methylguanine-DNA methyltransferase (MGMT): impact on cancer risk in response to tobacco smoke. *Mutation Research*. 2012; 736: 64–74.
- [15] Esteller M, Riques RA, Toyota M, Capella G, Moreno V, Peinado MA, *et al.* Promoter hypermethylation of the DNA repair gene O(6)-methylguanine-DNA methyltransferase is associated with the presence of G:C to A:T transition mutations in p53 in human colorectal tumorigenesis. *Cancer Research*. 2001; 61: 4689–4692.
- [16] Yuan X, Xu J, Fang W, Zhao Z, Wang F, Tong Z. The Association Between MGMT Promoter Methylation and Patients with Gastric Cancer: A Meta-Analysis. *Genetic Testing and Molecular Biomarkers*. 2017; 21: 213–221.
- [17] Kang B, Lee HS, Jeon SW, Park SY, Choi GS, Lee WK, *et al.* Progressive alteration of DNA methylation of Alu, MGMT, MINT2, and TFPI2 genes in colonic mucosa during colorectal cancer development. *Cancer Biomarkers*. 2021; 32: 231–236.
- [18] Paydar P, Asadikaram G, Nejad HZ, Akbari H, Abolhassani M, Moazed V, *et al.* Epigenetic modulation of BRCA-1 and MGMT genes, and histones H4 and H3 are associated with breast tumors. *Journal of Cellular Biochemistry*. 2019; 120: 13726–13736.
- [19] Sawhney M, Rohatgi N, Kaur J, Gupta SD, Deo SVS, Shukla NK, *et al.* MGMT expression in oral precancerous and cancerous lesions: correlation with progression, nodal metastasis and poor prognosis. *Oral Oncology*. 2007; 43: 515–522.
- [20] Zhang Z, Xin S, Gao M, Cai Y. Promoter hypermethylation of MGMT gene may contribute to the pathogenesis of gastric cancer: A PRISMA-compliant meta-analysis. *Medicine*. 2017; 96: e6708.
- [21] Chen YX, He LL, Xiang XP, Shen J, Qi HY. O⁶-methylguanine DNA methyltransferase is upregulated in malignant transformation of gastric epithelial cells *via* its gene promoter DNA hypomethylation. *World Journal of Gastrointestinal Oncology*. 2022; 14: 664–677.
- [22] Zhang YJ, Chen Y, Ahsan H, Lunn RM, Lee PH, Chen CJ, *et al.* Inactivation of the DNA repair gene O⁶-methylguanine-DNA methyltransferase by promoter hypermethylation and its relationship to aflatoxin B1-DNA adducts and p53 mutation in hepatocellular carcinoma. *International Journal of Cancer*. 2003; 103: 440–444.
- [23] Niger M, Nichetti F, Casadei-Gardini A, Morano F, Pircher C, Tamborini E, *et al.* MGMT inactivation as a new biomarker in patients with advanced biliary tract cancers. *Molecular Oncology*. 2022; 16: 2733–2746.
- [24] Yang Z, Li F. O-6-methylguanine-DNA methyltransferase gene promoter methylation and lung cancer risk: A meta-analysis. *Journal of Cancer Research and Therapeutics*. 2016; 12: C233–C236.
- [25] Martínez-Ramírez OC, Pérez-Morales R, Castro-Hernández C, Gonsébat ME, Casas-Ávila L, Valdés-Flores M, *et al.* Association of the Promoter Methylation and the rs12917 Polymorphism of MGMT with Formation of DNA Bulky Adducts and the Risk

of Lung Cancer in Mexican Mestizo Population. *DNA and Cell Biology*. 2019; 38: 307–313.

- [26] Pegg AE, Fang Q, Loktionova NA. Human variants of O6-alkylguanine-DNA alkyltransferase. *DNA Repair*. 2007; 6: 1071–1078.
- [27] Moitra P, Chatterjee A, Kota PK, Epari S, Patil V, Dasgupta A, *et al.* Temozolomide-induced myelotoxicity and single nucleotide polymorphisms in the MGMT gene in patients with adult diffuse glioma: a single-institutional pharmacogenetic study. *Journal of Neuro-Oncology*. 2022; 156: 625–634.
- [28] Kewitz-Hempel S, Kurch L, Cepelova M, Volkmer I, Sauerbrey A, Conrad E, *et al.* Impact of rs12917 MGMT Polymorphism on [(18)F]FDG-PET Response in Pediatric Hodgkin Lymphoma (PHL). *Molecular Imaging and Biology*. 2019; 21: 1182–1191.
- [29] Inoue R, Isono M, Abe M, Abe T, Kobayashi H. A genotype of the polymorphic DNA repair gene MGMT is associated with de novo glioblastoma. *Neurological Research*. 2003; 25: 875–879.
- [30] Sharma S, Salehi F, Scheithauer BW, Rotondo F, Syro LV, Kovacs K. Role of MGMT in tumor development, progression, diagnosis, treatment and prognosis. *Anticancer Research*. 2009; 29: 3759–3768.
- [31] Hegi ME, Diserens AC, Godard S, Dietrich PY, Regli L, Ostermann S, *et al.* Clinical trial substantiates the predictive value of O-6-methylguanine-DNA methyltransferase promoter methylation in glioblastoma patients treated with temozolomide. *Clinical Cancer Research*. 2004; 10: 1871–1874.
- [32] Hegi ME, Diserens AC, Gorlia T, Hamou MF, de Tribolet N, Weller M, *et al.* MGMT gene silencing and benefit from temozolomide in glioblastoma. *The New England Journal of Medicine*. 2005; 352: 997–1003.
- [33] Haque W, Teh C, Butler EB, Teh BS. Prognostic and predictive impact of MGMT promoter methylation status in high risk grade II glioma. *Journal of Neuro-Oncology*. 2022; 157: 137–146.
- [34] Haque W, Thong E, Andrabi S, Verma V, Brian Butler E, Teh BS. Prognostic and predictive impact of MGMT promoter methylation in grade 3 gliomas. *Journal of Clinical Neuroscience*. 2021; 85: 115–121.
- [35] Szyllberg M, Sokal P, Śledzińska P, Bebyn M, Krajewski S, Szyllberg L, *et al.* MGMT Promoter Methylation as a Prognostic Factor in Primary Glioblastoma: A Single-Institution Observational Study. *Biomedicines*. 2022; 10: 2030.
- [36] Costa BM, Caeiro C, Guimarães I, Martinho O, Jaraquemada T, Augusto I, *et al.* Prognostic value of MGMT promoter methylation in glioblastoma patients treated with temozolomide-based chemoradiation: a Portuguese multicentre study. *Oncology Reports*. 2010; 23: 1655–1662.
- [37] Egaña L, Auzmendi-Iriarte J, Andermatten J, Villanua J, Ruiz I, Elua-Pinin A, *et al.* Methylation of MGMT promoter does not predict response to temozolomide in patients with glioblastoma in Donostia Hospital. *Scientific Reports*. 2020; 10: 18445.
- [38] Dahlrot RH, Larsen P, Boldt HB, Kreutzfeldt MS, Hansen S, Hjelmborg JB, *et al.* Posttreatment Effect of MGMT Methylation Level on Glioblastoma Survival. *Journal of Neuropathology and Experimental Neurology*. 2019; 78: 633–640.
- [39] Caccese M, Simonelli M, Villani V, Rizzato S, Ius T, Pasqualetti F, *et al.* Definition of the Prognostic Role of MGMT Promoter Methylation Value by Pyrosequencing in Newly Diagnosed IDH Wild-Type Glioblastoma Patients Treated with Radiochemotherapy: A Large Multicenter Study. *Cancers*. 2022; 14: 2425.
- [40] Schulze Heuling E, Knab F, Radke J, Eskilsson E, Martinez-Ledesma E, Koch A, *et al.* Prognostic Relevance of Tumor Purity and Interaction with MGMT Methylation in Glioblastoma. *Molecular Cancer Research*. 2017; 15: 532–540.
- [41] Lou L, Zhang W, Li J, Wang Y. Abnormal MGMT Promoter Methylation in Gastrointestinal Stromal Tumors: Genetic Susceptibility and Association with Clinical Outcome. *Cancer Management and Research*. 2020; 12: 9941–9952.
- [42] Cros J, Hentic O, Rebours V, Zappa M, Gille N, Theou-Anton N, *et al.* MGMT expression predicts response to temozolomide in pancreatic neuroendocrine tumors. *Endocrine-related Cancer*. 2016; 23: 625–633.
- [43] Amatu A, Sartore-Bianchi A, Moutinho C, Belotti A, Bencardino K, Chirico G, *et al.* Promoter CpG island hypermethylation of the DNA repair enzyme MGMT predicts clinical response to dacarbazine in a phase II study for metastatic colorectal cancer. *Clinical Cancer Research*. 2013; 19: 2265–2272.
- [44] Tuominen R, Jewell R, van den Oord JJ, Wolter P, Stiermer U, Lindholm C, *et al.* MGMT promoter methylation is associated with temozolomide response and prolonged progression-free survival in disseminated cutaneous melanoma. *International Journal of Cancer*. 2015; 136: 2844–2853.
- [45] Toffolatti L, Scquizzato E, Cavallin S, Canal F, Scarpa M, Stefani PM, *et al.* MGMT promoter methylation and correlation with protein expression in primary central nervous system lymphoma. *Virchows Archiv*. 2014; 465: 579–586.
- [46] Rehman AU, Saikia S, Iqbal MA, Ahmad I, Sadaf, Anees A, *et al.* Decreased expression of MGMT in correlation with aberrant DNA methylation in esophageal cancer patients from North India. *Tumour Biology*. 2017; 39: 1010428317705770.
- [47] Chen R, Zheng Y, Zhuo L, Wang S. Association between MGMT Promoter Methylation and Risk of Breast and Gynecologic Cancers: A Systematic Review and Meta-Analysis. *Scientific Reports*. 2017; 7: 12783.
- [48] Fietkau R, Putz F, Lahmer G, Semrau S, Buslei R. Can MGMT promoter methylation status be used as a prognostic and predictive marker for glioblastoma multiforme at the present time? A word of caution. *Strahlentherapie und Onkologie*. 2013; 189: 993–995.
- [49] Lee CC, Kuo YC, Hu JM, Chang PK, Sun CA, Yang T, *et al.* *MTNR1B* polymorphisms with *CDKN2A* and *MGMT* methylation status are associated with poor prognosis of colorectal cancer in Taiwan. *World Journal of Gastroenterology*. 2021; 27: 5737–5752.
- [50] Christmann M, Verbeek B, Roos WP, Kaina B. O(6)-Methylguanine-DNA methyltransferase (MGMT) in normal tissues and tumors: enzyme activity, promoter methylation and immunohistochemistry. *Biochimica et Biophysica Acta*. 2011; 1816: 179–190.
- [51] Belanich M, Pastor M, Randall T, Guerra D, Kibitel J, Alas L, *et al.* Retrospective study of the correlation between the DNA repair protein alkyltransferase and survival of brain tumor patients treated with carmustine. *Cancer Research*. 1996; 56: 783–788.
- [52] Younis SG, Khedr RAEG, El-Shorbagy SH. Immunohistochemical analysis of O6-methylguanine-DNA methyltransferase (MGMT) protein expression as prognostic marker in glioblastoma patients treated with radiation therapy with concomitant and adjuvant Temozolomide. *Journal of the Egyptian National Cancer Institute*. 2016; 28: 23–30.
- [53] Butta S, Gupta MK. Immunohistochemical expression of MGMT in gliomas and its role in ascertaining patient survival. *Medicine and Pharmacy Reports*. 2021; 94: 318–324.
- [54] Ohno T, Hiraga J, Ohashi H, Sugisaki C, Li E, Asano H, *et al.* Loss of O6-methylguanine-DNA methyltransferase protein expression is a favorable prognostic marker in diffuse large B-cell lymphoma. *International Journal of Hematology*. 2006; 83: 341–347.
- [55] Mokhtar M, Kondo K, Namura T, Ali AHK, Fujita Y, Takai C, *et al.* Methylation and expression profiles of MGMT gene in thymic epithelial tumors. *Lung Cancer*. 2014; 83: 279–287.
- [56] Bengtsson D, Schröder HD, Andersen M, Maiter D, Berinder K, Feldt Rasmussen U, *et al.* Long-term outcome and MGMT as a predictive marker in 24 patients with atypical pituitary ade-

- nomas and pituitary carcinomas given treatment with temozolomide. *The Journal of Clinical Endocrinology and Metabolism*. 2015; 100: 1689–1698.
- [57] Kakkar A, Kumar A, Jha P, Goyal N, Mallick S, Sharma MC, *et al.* Meningeal hemangiopericytomas: a clinicopathological study with emphasis on MGMT (O(6)-methylguanine-DNA methyltransferase) promoter methylation status. *Neuropathology*. 2014; 34: 333–342.
- [58] Yang QC, Wang YH, Lin Y, Xue L, Chen YJ, Chen MH, *et al.* Expression of O(6)-methylguanine DNA methyltransferase (MGMT) and its clinical significance in gastroenteropancreatic neuroendocrine neoplasm. *International Journal of Clinical and Experimental Pathology*. 2014; 7: 4204–4212.
- [59] Schmitt AM, Pavel M, Rudolph T, Dawson H, Blank A, Komminoth P, *et al.* Prognostic and predictive roles of MGMT protein expression and promoter methylation in sporadic pancreatic neuroendocrine neoplasms. *Neuroendocrinology*. 2014; 100: 35–44.
- [60] Oliver JA, Ortiz R, Melguizo C, Alvarez PJ, Gómez-Millán J, Prados J. Prognostic impact of MGMT promoter methylation and MGMT and CD133 expression in colorectal adenocarcinoma. *BMC Cancer*. 2014; 14: 511.
- [61] Scesnaite A, Jarmalaite S, Mueller M, Agaimy A, Zenk J, Hartmann A, *et al.* Prognostic value of O-6-methylguanine-DNA methyltransferase loss in salivary gland carcinomas. *Head & Neck*. 2014; 36: 1258–1267.
- [62] Ban X, Mo S, Lu Z, Jia C, Shao H, Chang X, *et al.* Expression and methylation status of MMR and MGMT in well-differentiated pancreatic neuroendocrine tumors and potential clinical applications. *Endocrine*. 2022; 77: 538–545.
- [63] Mineura K, Izumi I, Watanabe K, Kowada M. O6-alkylguanine-DNA alkyltransferase activity in human brain tumors. *The Tohoku Journal of Experimental Medicine*. 1991; 165: 223–228.
- [64] Silber JR, Bobola MS, Ghatan S, Blank A, Kolstoe DD, Berger MS. O6-methylguanine-DNA methyltransferase activity in adult gliomas: relation to patient and tumor characteristics. *Cancer Research*. 1998; 58: 1068–1073.
- [65] Bobola MS, Alnoor M, Chen JYS, Kolstoe DD, Silbergeld DL, Rostomily RC, *et al.* O⁶-methylguanine-DNA methyltransferase activity is associated with response to alkylating agent therapy and with MGMT promoter methylation in glioblastoma and anaplastic glioma. *BBA Clinical*. 2015; 3: 1–10.
- [66] Pegg AE, Dolan ME, Moschel RC. Structure, function, and inhibition of O6-alkylguanine-DNA alkyltransferase. *Progress in Nucleic Acid Research and Molecular Biology*. 1995; 51: 167–223.
- [67] Gilbert MR, Wang M, Aldape KD, Stupp R, Hegi ME, Jaeckle KA, *et al.* Dose-dense temozolomide for newly diagnosed glioblastoma: a randomized phase III clinical trial. *Journal of Clinical Oncology*. 2013; 31: 4085–4091.
- [68] Geng X, Zhang Y, Lin X, Zeng Z, Hu J, Hao L, *et al.* Exosomal circWDR62 promotes temozolomide resistance and malignant progression through regulation of the miR-370-3p/MGMT axis in glioma. *Cell Death & Disease*. 2022; 13: 596.
- [69] Zhou Y, Chen L, Ding D, Li Z, Cheng L, You Q, *et al.* Cyanidin-3-O-glucoside inhibits the β -catenin/MGMT pathway by up-regulating miR-214-5p to reverse chemotherapy resistance in glioma cells. *Scientific Reports*. 2022; 12: 7773.
- [70] Cheng M, Wang Q, Chen L, Zhao D, Tang J, Xu J, *et al.* LncRNA UCA1/miR-182-5p/MGMT axis modulates glioma cell sensitivity to temozolomide through MGMT-related DNA damage pathways. *Human Pathology*. 2022; 123: 59–73.
- [71] Wu K, Wang Q, Liu YL, Xiang Z, Wang QQ, Yin L, *et al.* LncRNA POU3F3 Contributes to Dacarbazine Resistance of Human Melanoma Through the MiR-650/MGMT Axis. *Frontiers in Oncology*. 2021; 11: 643613.
- [72] Li X, Xu X, Chen K, Wu H, Wang Y, Yang S, *et al.* miR-370 Sensitizes TMZ Response Dependent of MGMT Status in Primary Central Nervous System Lymphoma. *Pathology Oncology Research*. 2020; 26: 707–714.
- [73] Wu X, Luo Q, Zhao P, Chang W, Wang Y, Shu T, *et al.* MGMT-activated DUB3 stabilizes MCL1 and drives chemoresistance in ovarian cancer. *Proceedings of the National Academy of Sciences of the United States of America*. 2019; 116: 2961–2966.
- [74] Esteller M, Toyota M, Sanchez-Cespedes M, Capella G, Peinado MA, Watkins DN, *et al.* Inactivation of the DNA repair gene O6-methylguanine-DNA methyltransferase by promoter hypermethylation is associated with G to A mutations in K-ras in colorectal tumorigenesis. *Cancer Research*. 2000; 60: 2368–2371.
- [75] Franco Pinto J, Fillion A, Duchambon P, Bombard S, Granzhan A. Acridine-O⁶-benzylguanine hybrids: Synthesis, DNA binding, MGMT inhibition and antiproliferative activity. *European Journal of Medicinal Chemistry*. 2022; 227: 113909.
- [76] Feng X, Xue F, He G, Ni Q, Huang S. Banxia xiexin decoction affects drug sensitivity in gastric cancer cells by regulating MGMT expression via IL 6/JAK/STAT3 mediated PD L1 activity. *International Journal of Molecular Medicine*. 2021; 48: 165.
- [77] Lei Y, Tang L, Hu J, Wang S, Liu Y, Yang M, *et al.* Inhibition of MGMT-mediated autophagy suppression decreases cisplatin chemosensitivity in gastric cancer. *Biomedicine & Pharmacotherapy*. 2020; 125: 109896.
- [78] Li Q, Ren B, Gui Q, Zhao J, Wu M, Shen M, *et al.* Blocking MAPK/ERK pathway sensitizes hepatocellular carcinoma cells to temozolomide via downregulating MGMT expression. *Annals of Translational Medicine*. 2020; 8: 1305.
- [79] Li X, Yang C, Luo N, Yang Y, Guo Y, Chen P, *et al.* Ubiquitination and degradation of MGMT by TRIM72 increases the sensitivity of uveal melanoma cells to Dacarbazine treatment. *Cancer Biomarkers*. 2022; 34: 275–284.
- [80] Jin L, Kiang KMY, Cheng SY, Leung GKK. Pharmacological inhibition of serine synthesis enhances temozolomide efficacy by decreasing O⁶-methylguanine DNA methyltransferase (MGMT) expression and reactive oxygen species (ROS)-mediated DNA damage in glioblastoma. *Laboratory Investigation*. 2022; 102: 194–203.
- [81] Tang Q, Cao H, Tong N, Liu Y, Wang W, Zou Y, *et al.* Tubeimoside-I sensitizes temozolomide-resistant glioblastoma cells to chemotherapy by reducing MGMT expression and suppressing EGFR induced PI3K/Akt/mTOR/NF- κ B-mediated signaling pathway. *Phytomedicine*. 2022; 99: 154016.
- [82] Liu Y, Du Z, Xu Z, Jin T, Xu K, Huang M, *et al.* Overexpressed GNA13 induces temozolomide sensitization via downregulating MGMT and p-RELA in glioma. *American Journal of Translational Research*. 2021; 13: 11413–11426.
- [83] Li H, Liu S, Jin R, Xu H, Li Y, Chen Y, *et al.* Pyrrvinium pamoate regulates MGMT expression through suppressing the Wnt/ β -catenin signaling pathway to enhance the glioblastoma sensitivity to temozolomide. *Cell Death Discovery*. 2021; 7: 288.
- [84] Shi J, Chen G, Dong X, Li H, Li S, Cheng S, *et al.* METTL3 Promotes the Resistance of Glioma to Temozolomide via Increasing MGMT and ANPG in a m⁶A Dependent Manner. *Frontiers in Oncology*. 2021; 11: 702983.
- [85] Lv W, Li Q, Jia B, He Y, Ru Y, Guo Q, *et al.* Differentiated embryonic chondrocyte-expressed gene 1 promotes temozolomide resistance by modulating the SP1-MGMT axis in glioblastoma. *American Journal of Translational Research*. 2021; 13: 2331–2349.
- [86] Lin K, Gueble SE, Sundaram RK, Huseman ED, Bindra RS, Herzon SB. Mechanism-based design of agents that selectively target drug-resistant glioma. *Science*. 2022; 377: 502–511.
- [87] Lim M, Weller M, Idhah A, Steinbach J, Finocchiaro G, Raval RR, *et al.* Phase III trial of chemoradiotherapy with

- temozolomide plus nivolumab or placebo for newly diagnosed glioblastoma with methylated MGMT promoter. *Neuro-Oncology*. 2022; 24: 1935–1949.
- [88] Omuro A, Brandes AA, Carpentier AF, Idhah A, Reardon DA, Cloughesy T, *et al.* Radiotherapy combined with nivolumab or temozolomide for newly diagnosed glioblastoma with unmethylated MGMT promoter: An international randomized phase III trial. *Neuro-Oncology*. 2023; 25: 123–134.
- [89] Weller M, Tabatabai G, Kästner B, Felsberg J, Steinbach JP, Wick A, *et al.* MGMT Promoter Methylation Is a Strong Prognostic Biomarker for Benefit from Dose-Intensified Temozolomide Rechallenge in Progressive Glioblastoma: The DIRECTOR Trial. *Clinical Cancer Research*. 2015; 21: 2057–2064.
- [90] Barazzuol L, Jena R, Burnet NG, Meira LB, Jeynes JCG, Kirkby KJ, *et al.* Evaluation of poly (ADP-ribose) polymerase inhibitor ABT-888 combined with radiotherapy and temozolomide in glioblastoma. *Radiation Oncology*. 2013; 8: 65.
- [91] Jue TR, Nozue K, Lester AJ, Joshi S, Schroder LBW, Whittaker SP, *et al.* Veliparib in combination with radiotherapy for the treatment of MGMT unmethylated glioblastoma. *Journal of Translational Medicine*. 2017; 15: 61.
- [92] Sim HW, McDonald KL, Lwin Z, Barnes EH, Rosenthal M, Foote MC, *et al.* A randomized phase II trial of veliparib, radiotherapy, and temozolomide in patients with unmethylated MGMT glioblastoma: the VERTU study. *Neuro-oncology*. 2021; 23: 1736–1749.
- [93] Schneider M, Potthoff AL, Evert BO, Dicks M, Ehrentauf D, Dolf A, *et al.* Inhibition of Intercellular Cytosolic Traffic via Gap Junctions Reinforces Lomustine-Induced Toxicity in Glioblastoma Independent of MGMT Promoter Methylation Status. *Pharmaceutics*. 2021; 14: 195.
- [94] Kirstein A, Schmid TE, Combs SE. The Role of miRNA for the Treatment of MGMT Unmethylated Glioblastoma Multiforme. *Cancers*. 2020; 12: 1099.
- [95] Katsigiannis S, Grau S, Krischek B, Er K, Pintea B, Goldbrunner R, *et al.* MGMT-Positive vs MGMT-Negative Patients With Glioblastoma: Identification of Prognostic Factors and Resection Threshold. *Neurosurgery*. 2021; 88: E323–E329.
- [96] Gessler F, Bernstock JD, Braczynski A, Lescher S, Baumgarten P, Harter PN, *et al.* Surgery for Glioblastoma in Light of Molecular Markers: Impact of Resection and MGMT Promoter Methylation in Newly Diagnosed IDH-1 Wild-Type Glioblastomas. *Neurosurgery*. 2019; 84: 190–197.
- [97] Stupp R, Mason WP, van den Bent MJ, Weller M, Fisher B, Taphoorn MJB, *et al.* Radiotherapy plus concomitant and adjuvant temozolomide for glioblastoma. *The New England Journal of Medicine*. 2005; 352: 987–996.
- [98] Curran WJ, Jr, Scott CB, Horton J, Nelson JS, Weinstein AS, Fischbach AJ, *et al.* Recursive partitioning analysis of prognostic factors in three Radiation Therapy Oncology Group malignant glioma trials. *Journal of the National Cancer Institute*. 1993; 85: 704–710.
- [99] Yuen CA, Barbaro M, Haggagi A. Newly Diagnosed Glioblastoma in Elderly Patients. *Current Oncology Reports*. 2022; 24: 325–334.
- [100] Wang Z, Du F, Ren Y, Jiang W. Treatment of MGMT promoter unmethylated glioblastoma with PD-1 inhibitor combined with anti-angiogenesis and epidermal growth factor receptor tyrosine kinase inhibitor: a case report. *Annals of Translational Medicine*. 2021; 9: 1508.
- [101] Yu W, Zhang L, Wei Q, Shao A. O⁶-Methylguanine-DNA Methyltransferase (MGMT): Challenges and New Opportunities in Glioma Chemotherapy. *Frontiers in Oncology*. 2020; 9: 1547.
- [102] Yang P, Zhang W, Wang Y, Peng X, Chen B, Qiu X, *et al.* IDH mutation and MGMT promoter methylation in glioblastoma: results of a prospective registry. *Oncotarget*. 2015; 6: 40896–40906.
- [103] Chai R, Li G, Liu Y, Zhang K, Zhao Z, Wu F, *et al.* Predictive value of MGMT promoter methylation on the survival of TMZ treated IDH-mutant glioblastoma. *Cancer Biology & Medicine*. 2021; 18: 272–282.
- [104] Nguyen HN, Lie A, Li T, Chowdhury R, Liu F, Ozer B, *et al.* Human TERT promoter mutation enables survival advantage from MGMT promoter methylation in IDH1 wild-type primary glioblastoma treated by standard chemoradiotherapy. *Neuro-Oncology*. 2017; 19: 394–404.
- [105] Vuong HG, Nguyen TQ, Ngo TNM, Nguyen HC, Fung KM, Dunn IF. The interaction between TERT promoter mutation and MGMT promoter methylation on overall survival of glioma patients: a meta-analysis. *BMC Cancer*. 2020; 20: 897.
- [106] Tunthanathip T, Sangkhathat S, Tanvejsilp P, Kanjanapradit K. Prognostic Impact of the Combination of MGMT Methylation and TERT Promoter Mutation in Glioblastoma. *Journal of Neurosciences in Rural Practice*. 2021; 12: 694–703.
- [107] Weller M, van den Bent M, Preusser M, Le Rhun E, Tonn JC, Minniti G, *et al.* EANO guidelines on the diagnosis and treatment of diffuse gliomas of adulthood. *Nature Reviews. Clinical Oncology*. 2021; 18: 170–186.
- [108] Sasai K, Nodagashira M, Nishihara H, Aoyanagi E, Wang L, Katoh M, *et al.* Careful exclusion of non-neoplastic brain components is required for an appropriate evaluation of O⁶-methylguanine-DNA methyltransferase status in glioma: relationship between immunohistochemistry and methylation analysis. *The American Journal of Surgical Pathology*. 2008; 32: 1220–1227.
- [109] Quillien V, Lavenue A, Ducray F, Joly MO, Chinot O, Fina F, *et al.* Validation of the high-performance of pyrosequencing for clinical MGMT testing on a cohort of glioblastoma patients from a prospective dedicated multicentric trial. *Oncotarget*. 2016; 7: 61916–61929.
- [110] Xie H, Tubbs R, Yang B. Detection of MGMT promoter methylation in glioblastoma using pyrosequencing. *International Journal of Clinical and Experimental Pathology*. 2015; 8: 1790–1796.
- [111] Nguyen N, Redfield J, Ballo M, Michael M, Sorenson J, Dibaba D, *et al.* Identifying the optimal cutoff point for MGMT promoter methylation status in glioblastoma. *CNS Oncology*. 2021; 10: CNS74.
- [112] Hosoya T, Takahashi M, Honda-Kitahara M, Miyakita Y, Ohno M, Yanagisawa S, *et al.* MGMT gene promoter methylation by pyrosequencing method correlates volumetric response and neurological status in IDH wild-type glioblastomas. *Journal of Neuro-Oncology*. 2022; 157: 561–571.
- [113] Jahin M, Fenech-Salerno B, Moser N, Georgiou P, Flanagan J, Toumazou C, *et al.* Detection of MGMT methylation status using a Lab-on-Chip compatible isothermal amplification method. *Annual International Conference of the IEEE Engineering in Medicine and Biology Society. IEEE Engineering in Medicine and Biology Society. Annual International Conference*. 2021; 2021: 7385–7389.
- [114] Rosas-Alonso R, Colmenarejo-Fernandez J, Pernia O, Rodriguez-Antolín C, Esteban I, Ghanem I, *et al.* Clinical validation of a novel quantitative assay for the detection of MGMT methylation in glioblastoma patients. *Clinical Epigenetics*. 2021; 13: 52.
- [115] von Rosenstiel C, Wiestler B, Haller B, Schmidt-Graf F, Gempt J, Bettstetter M, *et al.* Correlation of the quantitative level of MGMT promoter methylation and overall survival in primary diagnosed glioblastomas using the quantitative MethyQESD method. *Journal of Clinical Pathology*. 2020; 73: 112–115.
- [116] Sahara N, Hartanto RA, Yoshuantari N, Dananjoyo K, Widodo I, Malueka RG, *et al.* Diagnostic Accuracy of Immunohisto-

chemistry in Detecting MGMT Methylation Status in Patients with Glioma. *Asian Pacific Journal of Cancer Prevention*. 2021; 22: 3803–3808.

- [117] Hegi ME, Genbrugge E, Gorlia T, Stupp R, Gilbert MR, Chinot OL, *et al.* MGMT Promoter Methylation Cutoff with Safety Margin for Selecting Glioblastoma Patients into Trials Omitting Temozolomide: A Pooled Analysis of Four Clinical Trials. *Clinical Cancer Research*. 2019; 25: 1809–1816.
- [118] Li M, Dong G, Zhang W, Ren X, Jiang H, Yang C, *et al.* Combining MGMT promoter pyrosequencing and protein expression to optimize prognosis stratification in glioblastoma. *Cancer Science*. 2021; 112: 3699–3710.
- [119] McGarry SD, Hurrell SL, Kaczmarowski AL, Cochran EJ, Connelly J, Rand SD, *et al.* Magnetic Resonance Imaging-Based Radiomic Profiles Predict Patient Prognosis in Newly Diagnosed Glioblastoma Before Therapy. *Tomography*. 2016; 2: 223–228.
- [120] Salihoğlu YS, Uslu Erdemir R, Aydur Püren B, Özdemir S, Uyulan Ç, Ergüzel TT, *et al.* Diagnostic Performance of Machine Learning Models Based on ¹⁸F-FDG PET/CT Radiomic Features in the Classification of Solitary Pulmonary Nodules. *Molecular Imaging and Radionuclide Therapy*. 2022; 31: 82–88.
- [121] Le NQK, Do DT, Chiu FY, Yapp EKY, Yeh HY, Chen CY. XGBoost Improves Classification of MGMT Promoter Methylation Status in IDH1 Wildtype Glioblastoma. *Journal of Personalized Medicine*. 2020; 10: 128.
- [122] Do DT, Yang MR, Lam LHT, Le NQK, Wu YW. Improving MGMT methylation status prediction of glioblastoma through optimizing radiomics features using genetic algorithm-based machine learning approach. *Scientific Reports*. 2022; 12: 13412.
- [123] Lu J, Li X, Li H. Perfusion parameters derived from MRI for preoperative prediction of IDH mutation and MGMT promoter methylation status in glioblastomas. *Magnetic Resonance Imaging*. 2021; 83: 189–195.
- [124] Yogananda CGB, Shah BR, Nalawade SS, Murugesan GK, Yu FF, Pinho MC, *et al.* MRI-Based Deep-Learning Method for Determining Glioma MGMT Promoter Methylation Status. *American Journal of Neuroradiology*. 2021; 42: 845–852.
- [125] Weaver KD, Grossman SA, Herman JG. Methylated tumor-specific DNA as a plasma biomarker in patients with glioma. *Cancer Investigation*. 2006; 24: 35–40.
- [126] Fiano V, Trevisan M, Trevisan E, Senetta R, Castiglione A, Sacerdote C, *et al.* MGMT promoter methylation in plasma of glioma patients receiving temozolomide. *Journal of Neuro-Oncology*. 2014; 117: 347–357.