Check for updates

Value of Thallium-201 SPECT and SPECT/CT Brain Imaging in Differentiating Malignant From Nonmalignant Lesions: A Comparative Case-Series Study With Pathologic and/or Clinical Correlation

Nicole Girlyn T. Pang^{1*}, Julette Marie F. Batara^{1,2*}, Lara Triccia C. Luistro³, Christianne V. Mojica¹, Allan Jay C. Domingo³

¹Institute for Neurosciences, St. Luke's Medical Center-Quezon City and Global City, Philippines ²Department of Neurosciences, College of Medicine and Philippine General Hospital, University of the Philippines Manila, Manila, Philippines ³Department of Nuclear Medicine, St. Luke' Medical Center-Global City, Taguig, Philippines

Received June 11, 2023 Revised July 25, 2023 Accepted August 21, 2023

Correspondence

Nicole Girlyn T Pang Institute for Neurosciences, St. Luke's Medical Center Quezon City, 279 E. Rodriguez Sr. Ave., Quezon City, 1112, Metro Manila, Philippines Tel: +63-917-5677264 E-mail: ngtpang@gmail.com

Julette Marie F. Batara Institute for Neurosciences. St. Luke's Medical Center Quezon City. 279 E. Rodriguez Sr. Ave., Quezon City, 1112, Metro Manila, Philippines Tel: +63-917-8662388 E-mail: jbatara1@yahoo.com

*Nicole Girlyn T. Pang and Julette Marie F. Batara contributed equally to this work.

Background Thallium-201 single-photon emission computed tomography/computed tomography (SPECT/CT) brain scan is an imaging modality which can be done to differentiate between malignant and nonmalignant lesions among patients with nonconclusive findings on conventional neuroimaging. This study describes the results of thallium-201 SPECT/CT brain imaging and relate it to histopathologic and/or clinical findings and evaluate the value of thallium-201 SPECT/CT brain imaging in differentiating malignant from nonmalignant lesions.

Methods This is a retrospective case series study of 10 patients with cerebral lesions who underwent thallium-201 SPECT/CT brain imaging in a hospital in the Philippines from 2010 to 2021.

A total of 10 patients underwent thallium-201 SPECT/CT brain scan. Six had nega-Results tive results while 4 had positive results. All of the patients who had positive results were found to have malignancy, whether recurrent or newly diagnosed. All of the patients with negative scan were found to have either an infectious and inflammatory disease and responded to treatment albeit in different degrees. Two of the 10 patients underwent biopsy whose results were consistent with the thallium-201 SPECT/CT brain scan results.

Thallium-201 brain scan combined with SPECT and SPECT/CT has been demon-Conclusion strated to be useful in distinguishing malignant from nonmalignant lesions and is more cost-effective versus other imaging techniques. The findings in this study support the role of thallium scintigraphy in the diagnosis of patients with brain lesions most significantly when there is a need to differentiate between a malignant and benign condition.

Keywords Thallium; Tomography, emission-computed, single-photon; Neuroimaging; Brain tumor; Malignancy.

INTRODUCTION

The primary neuroimaging methods for the identification

and localization of intracranial lesions are MRI and CT neuroimaging. Of the two, MRI is the imaging modality of choice to evaluate brain tumor location, size, and extent, mass effect, and involvement of critical structures [1]. However, there are instances where MRI cannot accurately determine characteristics of lesions to differentiate malignant lesions from nonmalignant ones. Primary central nervous system (CNS) lymphoma and various CNS infections may have indistinguishable

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (https://creativecommons.org/licenses/by-nc/4.0) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

Copyright © 2023 The Korean Brain Tumor Society, The Korean Society for Neuro-Oncology, and The Korean Society for Pediatric Neuro-Oncology

characteristics of lesions on neuroimaging studies [2].

Brain biopsy remains to be the gold standard in the diagnosis of brain tumors, which is most often done at the time of tumor removal. Limitations on the use of brain biopsy include the hard-to-reach location of the tumor, patients who are poor surgical candidates, and when patient is not amenable to an invasive surgical procedure [3]. There are also instances such as in HIV patients particularly, it would be helpful in distinguishing parasitic infections from malignancies such as lymphoma where delaying treatment for a therapeutic trial would be detrimental.

Thallium-201 single-photon emission CT/CT (SPECT/CT) brain imaging has been reported to be more sensitive and specific in identifying carcinomatous cerebral lesions as compared to MRI, CT, or radionuclide studies using other radiotracers [4-9]. Separating malignant from benign lesions with thallium brain scans is based on the tracer activity imaged during the procedure. The normal absence of tracer uptake in the brain permits easier visualization of tumors and imaging of brain lesions has been the most common application of thallium in tumor imaging. There is differential washout of the tracer from malignant tumors versus normal brain tissues, with prolonged clearance in the former. Lymphoma has a moderate to intense tracer uptake while a CNS infection such as toxoplasmosis has absent uptake.

The aim of this retrospective study is to describe the results of thallium-201 SPECT/CT brain imaging and relate it to histopathologic and/or clinical findings and to evaluate the value of thallium-201 SPECT/CT brain imaging in differentiating malignant from nonmalignant lesions. This is the first study in the Philippines investigating the usefulness of thallium-201 SPECT/CT brain imaging.

MATERIALS AND METHODS

This is a retrospective case series of patients ages 18 years old and above with cerebral lesions who underwent thallium-201 SPECT/CT brain imaging at St. Luke's Medical Center–Global City, Philippines from 2010 to 2021. Patients who were not able to complete the thallium-201 brain imaging were excluded in the study.

Brain scintigraphy was performed using 4 mCi ±5% of thallium-201. Scanning was done using Siemens Symbia Intevo Bold or Siemens Symbia S dual-head hybrid SPECT/CT gamma camera (Siemens Healthineers, Erlangen, Germany) with a large field of view, low energy all purpose-parallel hole collimator. Images were acquired in continuous mode, non-circular orbit, 180 degrees, 32 views, 50 seconds per view with 128×128 matrix size. Initial planar images of the head were acquired 15 minutes post-intravenous injection of the radiotracer. Delayed planar images were acquired 90 minutes post-tracer injection. SPECT imaging of the head was done after the delayed scan and SPECT/CT was performed as needed. A positive thallium-201 SPECT brain imaging was interpreted based on the tracer uptake on the area of interest. All imaging findings were interpreted and reviewed by experienced nuclear medicine physicians.

The clinical protocol and all relevant documents were approved by the Institutional Ethics Review Committee. Each patient document was coded and does not contain any identifying information to ensure confidentiality.

The list of patients who underwent thallium-201 SPECT brain imaging were retrieved from the computerized database/ online imaging portal of the hospital. Data were gathered from review of medical records and healthcare system. Data pertaining to the outcomes of patients who underwent thallium 201 SPECT/CT brain imaging were recorded in the researchers' database. This included patients' demographic profile, comorbidities, neurologic symptoms, brain scintigraphy result, other imaging/ancillary results, histopathologic findings and clinical diagnosis, management, interventions and clinical outcome.

Descriptive statistics were used to summarize the demographic and clinical characteristics of the patients. The thallium-201 SPECT/CT brain imaging findings were compared to the final diagnosis of the patients and to histopathologic findings, if available.

Chart review was done for all the patients included in this study. No direct contact was done between the researchers and the patients. Identifying data of the patients have not been included in the data collection form to maintain confidentiality. This study with reference number SL-21298 received ethical clearance from the Institutional Ethics Review Committee of St. Luke's Medical Center on October 2021. Informed consent was wavied by the Institutional Review Committee.

RESULTS

The characteristics of patients who underwent thallium-201 brain imaging are summarized in Table 1. Half of the population belong to the 41 to 50-year age group. Most of the patients (80%) were females. Three of the 10 patients had a known carcinoma. The most common complaint of the patients tested was headache.

Four patients rendered positive results with thallium-201 brain scan while the rest have negative results. Of patients with positive thallium-201 brain scan, 3 patients were diagnosed with brain metastasis while 1 patient was diagnosed with pilocytic astrocytoma. Among patients with negative thallium-201 brain scan results, 5 patients were diagnosed with CNS infec-

le 1. C lient 1 1 1 6 6 6 6 7 7 7 7	haract Age (yr) 48 48 59 59 59 59 29 41 41 49	eristics of Sex Male Female Female Female Female Female	patients who underwee Comorbidities Meningioma, post-resection Lung cancer Hypertension, diabetes mellitus Cerebral toxoplasmosis Breast cancer None Cervical cancer	ent thallium-201 brain imaging Neurologic manifestations Asymptomatic Headache Blurring of vision Headache; perseveration; unilateral weakness Dizziness, vomiting, diplopia Dizziness Memory lapses; change in sensorium	Thallium-201 SPECT/CT Negative Negative Negative Positive Negative Positive	Histopathology - - Chronic inflammation and gliosis - -	Diagnosis Tuberculous granuloma Brain metastasis Neurocysticercosis Neurocysticercosis Cerebral toxoplasmosis recurrence Brain metastasis Tuberculous granuloma Brain metastasis	Outcome Responded to treatment Responded to treatment No response to treatment Responded to treatment Responded to treatment Responded to treatment Refused treatment
	48	Female	Hypertension	Headache; multiple craniopathies	Negative	•	Neurosarcoidosis	Responded to treatment
	22 48	Female Female	None Hypertension	Headache; dizziness Headache; unilateral weakness	Positive Negative	Pilocytic astrocytoma -	Astrocytoma Tuberculous granuloma	Responded to treatment Responded to treatment
CT/C1	l, single	e-photon (emission computed ton	nography/computed tomography				

NGT Pang et al.

tion while 1 patient was diagnosed with an autoimmune disease.

Patients were treated according to their final diagnosis. Six patients had good response to treatment, 1 patient is still ongoing treatment, 1 patient had poor response to treatment, and 1 patient refused treatment. Patient 2 died while still initiating treatment.

Only two patients consented and underwent biopsy of the brain lesion. Patient 3 who had negative result in thallium-201 brain scan underwent excision of cerebellar mass showed chronic inflammation and gliosis on biopsy and was diagnosed with neurocysticercosis. Patient 9 who had positive thallium-201 brain scan revealed pilocytic astrocytoma on biopsy and was treated accordingly. Figs. 1 and 2 show the thallium-201 brain scan findings and the corresponding histopathologic images of the biopsied cerebral lesions confirming the diagnosis. The thallium SPECT/CT scan images of the rest of the patients are shown in Supplementary Figs. 1 and 2 (in the online-only Data Supplement).

DISCUSSION

Neurologists may be faced with a diagnostic dilemma when a patient's clinical and radiographic presentation may point to several possible etiologies. As physicians, coming up with an accurate diagnosis is very crucial because treatment may vary depending on the disease entity and delay in the diagnosis may consequentially worsen a patient's condition. In case where our primary neuroimaging does not provide us with a definite diagnosis and brain biopsy may not be an option in some patients, thallium-201 brain scan may be used to supplement the diagnosis.

In our case series, these patients had nonconclusive primary neuroimaging findings which made their neurologists proceed with thallium-201 brain imaging to further support the diagnosis. Thallium-201 brain scan was able to accurately validate whether the patients have malignant lesions versus non-malignant lesions such as infection and inflammation. This aided the physicians on the treatment plan for the patients, on which they responded albeit in different degrees.

One of the advantages of thallium-201 SPECT/CT brain imaging in comparison with other imaging modality is its mechanism in localizing malignant lesions [10]. Thallium-201 uptake in the brain is influenced by regional blood flow, blood-brain barrier (BBB) permeability, cellular metabolic activity, and the variability in Na⁺-K⁺-ATPase (adenosine triphophatase) activity [11-14].

The integrity of the BBB makes it possible for different radiopharmaceuticals to distinguish between lesions that are benign and malignant. Non-malignant entities incite inflammatory responses that interrupt with physiological brain func-



Fig. 1. Thallium-201 brain scan findings and histopathologic image of Patient 3. A and B: A negative thallium brain scintigraphy planar (A) and SPECT/CT images (B) showing no increased tracer accumulation. C: Cerebellar tissues with gliosis (H&E, ×10). SPECT/CT, single-photon emission computed tomography/computed tomography.

tions which eventually lead to their clinical presentations. These pathologies are met by an immune response which does not cause a disruption in the BBB, limiting them to their area of origin and minimizing deviations from the body's homeostasis as much as possible. Brain tumors, on the other hand, encroach on the supply of normal tissues for oxygen and nutritional sustenance. They secrete angiogenesis factors to support their rapid growth. Blood vessels formed in the tumor-host interface do not have tight junctions, which normally only allow small amounts of plasma filtered at a time. This leakiness allows for tumor metastasis and serves as the entry point for thallium-201. It would follow that malignant processes would present as thallium-avid lesions in the brain and benign processes do not get visualized. It is also good to note that increase in BBB permeability alone will not necessarily increase thallium-201 uptake in the brain as some nonmalignant lesions may have increase in BBB permeability but will still only have little or no uptake [10].

Variability in thallium-201 uptake may as well provide further information on the brain lesion being investigated. The



Fig. 2. Thallium-201 brain scan findings and histopathologic image of Patient 9. A: Initial planar images after 90 minutes of thallium-201 intravenous injection showing no focus of increased tracer uptake. B: SPECT/CT imaging with a focus of thallium-201 uptake in the cervicomedullary junction and right brachium pontis. C: Cellular portion of brain tissue populated by bland round and spindle cells (H&E ×100), immunohistochemistry results revealed positive GFAP and P16, focally positive SOX10, IDH1 negative, and retained nuclear staining on ATRX favoring the diagnosis of pilocytic astrocytoma. SPECT/CT, single-photon emission computed tomography/computed tomography.

early uptake is thought to be related to tumor vascularity and the disruption of the BBB. Delayed uptake depends on the metabolic activity of the Na+-K+-ATPase mechanism of transmembrane transport into viable tumor cells, which may also indicate the degree of tumor malignancy. Thallium uptake is found to be significantly increased in tumors expressing increased vascular endothelial growth factor (VEGF) compared to VEGF weakly positive meningiomas in both the early image and the delayed image [15]. These mechanisms explain how malignant lesions have higher tracer activity in thallium brain scans compared to benign pathologies. By evaluating early and delayed uptake with normal "background" activity in thallium brain scan with SPECT, researchers have also come up with a retention index which correlates with degree of aggressiveness in gliomas and even meningiomas, Ki-67 proliferative index and predicting progression-free survival which are helpful for pretreatment planning. The addition of SPECT or SPECT/CT to the thallium-201 brain scan planar images provides improved sensitivity resulting in improved detection and localization.

Though in this case series we did not have a patient with a

case of radiation-induced necrosis, several studies reported use of thallium-201 SPECT/CT imaging in differentiating tumor recurrence from radiation-induced necrosis [14,16-18]. These two outcomes are most commonly encountered postbrain tumor treatment [19]. Their radiographic characteristics are similar and hard to distinguish on the conventional imaging which led to the use of several functional and physiologic imaging techniques to differentiate the two. A combination of MR spectroscopy and MR perfusion is commonly used but one of its disadvantages is its low accuracy when testing patients with coexisting tumor and necrosis [14]. In cases of malignant tumor recurrence, SPECT imaging will present with thallium-201 tracer uptake while radiation-induced necrosis will have no tracer uptake. In this case series, only Patients 1 and 2 had a history of cerebral lesion. Patient 1 had a resected meningioma and was found out to have a new cerebral lesion near the resected lesion during a surveillance MRI. No tracer uptake was seen on thallium-201 SPECT imaging thus new malignancy was ruled out. She was clinically diagnosed with tuberculous granuloma granuloma to which she responded to treatment with resolution of the previously

noted cerebral lesion on repeat imaging posttreatment. Patient 2 was previously treated for cerebral toxoplasmosis then developed new-onset neurologic deficits. MRI showed new multiple scattered cerebral lesions that had no tracer uptake on thallium brain scan. Patient was managed for cerebral toxoplasmosis recurrence and responded to treatment with resolution of neurologic deficits and cerebral lesions.

The sensitivity and specificity of thallium-201 SPECT in detecting and differentiating brain tumors was evaluated by Dierckx et al. [20] which resulted to 71.7% and 80.9%, respectively. Similar results were seen in the study of Hussain and Hussain [6] on the clinical utility of thallium-201 SPECT alone in the diagnosis of AIDS-related primary CNS lymphoma which yielded a sensitivity of 77% and specificity of 81%, but the diagnostic accuracy improved to 100% when it was used together with the cerebrospinal fluid Epstein-Barr virus polymerase chain reaction results. In a different study done by Lorberboym et al. [5], thallium-201 brain SPECT had a higher overall sensitivity of 100% and specificity of 90% in differentiating primary CNS lymphoma from cerebral toxoplasmosis. MR spectroscopy, in comparison, has a sensitivity of 89.3% and sensitivity of 73.9% [21]. Given these values, thallium-201 brain scan can competitively be used as an imaging option when other imaging modality is not convenient.

Thallium-201 brain scan is also a more cost-effective option among the noninvasive imaging techniques. As can be seen in Table 2, the gross price of a thallium-201 brain scan combined with SPECT/CT just a little above the price of a single cranial MRI with contrast and costs 35% less than an MR spectroscopy with perfusion study. In the Philippine setting where cost can be an important factor in the choice of diagnostic procedure, this study further supports the use of thallium-201 SPECT/CT scan in the diagnosis of patients with brain lesions.

The limitation of this study is the small number of patients included in the study and only 2 of the patients underwent biopsy to confirm the diagnosis. Nonetheless, objective clinical findings were used for the patients with no histopathologic results. A prospective study involving a larger population of sample with correlation of histopathologic findings is recommended to extensively examine the utility of thallium-201

 Table 2. Diagnostic neuroimaging procedure prices in a hospital in the Philippines

Diagnostic procedure	Prices (USD)
Thallium-201 brain scan	\$432.50
Thallium-201 brain scan + SPECT/CT	\$490.00
Cranial MRI with contrast	\$440.00
Cranial MRI with MRS and MRP	\$763.50

SPECT/CT, single-photon emission computed tomography/computed tomography; MRS, MR spectroscopy; MRP, MR perfusion brain scan.

In conclusion, thallium-201 brain scan combined with SPECT and SPECT/CT has been demonstrated to be useful in determining the nature of brain lesions, capable of distinguishing malignant process from nonmalignant ones, and more costeffective than other imaging techniques. Thallium-201 brain scan is a promising imaging modality to confirm diagnosis among patients with nonconclusive findings on conventional neuroimaging.

Supplementary Materials

The online-only Data Supplement is available with this article at https://doi.org/10.14791/btrt.2023.0022.

Availability of Data and Material

This database of the patients is under the ownership of the St. Luke's Medical Center - Global City. Inquiries regarding the access can be directed to the corresponding author.

ORCID iDs

Nicole Girlyn T. Pang Julette Marie F. Batara Lara Triccia C. Luistro Christianne V. Mojica Allan Jay C. Domingo https://orcid.org/0000-0001-5998-628X https://orcid.org/0000-0003-2824-4544 https://orcid.org/0000-0001-7475-5973 https://orcid.org/0000-0001-6628-1480 https://orcid.org/0000-0001-8079-9526

Author Contributions

Conceptualization: Nicole Girlyn T. Pang, Jullete Marie F. Batara. Data curation: Nicole Girlyn T. Pang, Jullete Marie F. Batara, Lara Triccia C. Luistro, Christianne V. Mojica. Formal analysis: Nicole Girlyn T. Pang, Lara Triccia C. Luistro, Christianne V. Mojica. Investigation: Nicole Girlyn T. Pang, Lara Triccia C. Luistro, Christianne V. Mojica. Methodology: Nicole Girlyn T. Pang, Jullete Marie F. Batara. Project administration: Nicole Girlyn T. Pang, Jullete Marie F. Batara. Resources: all authors. Software: Nicole Girlyn T. Pang, Lara Triccia C. Luistro. Supervision: Jullete Marie F. Batara, Allan Jay C. Domingo. Validation: all authors. Visualization: Jullete Marie F. Batara. Writing—original draft: Nicole Girlyn T. Pang, Jullete Marie F. Batara, Lara Triccia C. Luistro, Christianne V. Mojica. Writing–review & editing: Nicole Girlyn T. Pang, Jullete Marie F. Batara.

Conflicts of Interest

The authors have no potential conflicts of interest to disclose.

Funding Statement

None

Acknowledgments

We would like to thank our mentors, family, and friends for the motivation, guidance, and support in accomplishing this paper.

REFERENCES

- 1. Norden AD, Pope WB, Chang SM. Current concepts in brain tumor imaging. Am Soc Clin Oncol Educ Book 2012;32:119-24.
- Smego RA Jr, Orlovic D, Wadula J. An algorithmic approach to intracranial mass lesions in HIV/AIDS. Int J STD AIDS 2006;17:271-6.
- Di Bonaventura R, Montano N, Giordano M, Gessi M, Gaudino S, Izzo A, et al. Reassessing the role of brain tumor biopsy in the era of advanced surgical, molecular, and imaging techniques—a single-center experience with long-term follow-up. J Pers Med 2021;11:909.

- Lorberboym M, Estok L, Machac J, Germano I, Sacher M. Thallium-201 brain spect and CT/MRI imaging for differential diagnosis of cerebral toxoplasmosis and primary CNS lymphoma. Clin Nucl Med 1996;21:353.
- Lorberboym M, Estok L, Machac J, Germano I, Sacher M, Feldman R, et al. Rapid differential diagnosis of cerebral toxoplasmosis and primary central nervous system lymphoma by thallium-201 SPECT. J Nucl Med 1996;37:1150-4.
- Hussain FS, Hussain NS. Clinical utility of thallium-201 single photon emission computed tomography and cerebrospinal fluid Epstein-Barr virus detection using polymerase chain reaction in the diagnosis of AIDS-related primary central nervous system lymphoma. Cureus 2016;8:e606.
- Yang M, Sun J, Bai HX, Tao Y, Tang X, States LJ, et al. Diagnostic accuracy of SPECT, PET, and MRS for primary central nervous system lymphoma in HIV patients: a systematic review and meta-analysis. Medicine (Baltimore) 2017;96:e6676.
- Kaplan WD, Takvorian T, Morris JH, Rumbaugh CL, Connolly BT, Atkins HL. Thallium-201 brain tumor imaging: a comparative study with pathologic correlation. J Nucl Med 1987;28:47-52.
- Lorberboym M, Baram J, Feibel M, Hercbergs A, Lieberman L. A prospective evaluation of thallium-201 single photon emission computerized tomography for brain tumor burden. Int J Radiat Oncol Biol Phys 1995;32:249-54.
- Kim KT, Black KL, Marciano D, Mazziotta JC, Guze BH, Grafton S, et al. Thallium-201 SPECT imaging of brain tumors: methods and results. J Nucl Med 1990;31:965-9.
- 11. Atkins HL, Budinger TF, Lebowitz E, Ansari AN, Greene MW, Fairchild RG, et al. Thallium-201 for medical use. Part 3: human distribution and physical imaging properties. J Nucl Med 1977;18:133-40.
- 12. Bénard F, Romsa J, Hustinx R. Imaging gliomas with positron emission tomography and single-photon emission computed tomography.

Semin Nucl Med 2003;33:148-62.

- Kline JL, Noto RB, Glantz M. Single-photon emission CT in the evaluation of recurrent brain tumor in patients treated with gamma knife radiosurgery or conventional radiation therapy. AJNR Am J Neuroradiol 1996;17:1681-6.
- Tie J, Gunawardana DH, Rosenthal MA. Differentiation of tumor recurrence from radiation necrosis in high-grade gliomas using 201Tl-SPECT. J Clin Neurosci 2008;15:1327-34.
- Valotassiou V, Leondi A, Angelidis G, Psimadas D, Georgoulias P. SPECT and PET imaging of meningiomas. ScientificWorldJournal 2012;2012:412580.
- Matsunaga S, Shuto T, Takase H, Ohtake M, Tomura N, Tanaka T, et al. Semiquantitative analysis using thallium-201 SPECT for differential diagnosis between tumor recurrence and radiation necrosis after gamma knife surgery for malignant brain tumors. Int J Radiat Oncol Biol Phys 2013;85:47-52.
- Smith J, English J, Gilmore D. Evaluation of PET/CT and thallium-201 imaging for brain tumors vs. necrosis. J Nucl Med 2010;51(supplement 2):2120.
- Chochrek C, Stokes M, Gilmore D. Differentiating recurrent brain tumors from radiation necrosis with the use of thallium brain imaging. J Nucl Med 2009;50(supplement 2):2207.
- Verma N, Cowperthwaite MC, Burnett MG, Markey MK. Differentiating tumor recurrence from treatment necrosis: a review of neuro-oncologic imaging strategies. Neuro Oncol 2013;15:515-34.
- Dierckx RA, Martin JJ, Dobbeleir A, Crols R, Neetens I, De Deyn PP. Sensitivity and specificity of thallium-201 single-photon emission tomography in the functional detection and differential diagnosis of brain tumours. Eur J Nucl Med 1994;21:621-33.
- Munir S, Khan SA, Hanif H, Khan M. Diagnostic accuracy of magnetic resonance imaging in detection of intra-axial gliomas. Pak J Med Sci 2021;37:125-30.



Supplementary Fig. 1. Positive thallium-201 SPECT/CT brain scan. A: Patient 2: SPECT/CT imaging showing multiple foci of thallium tracer uptake in the brain. B: Patient 5: Thallium-201 brain scan SPECT fused with MRI showing uptake in the pontocerebellar area. C: Patient 7: Thallium SPECT/CT imaging showing multiple cerebral lesions with the most intense thallium tracer uptake on the left temporal lobe. D: Patient 9: SPECT/CT imaging with a focus of thallium-201 uptake in the cervicomedullary junction and right brachium pontis. SPECT/CT, single-photon emission computed tomography/computed tomography.



Supplementary Fig. 2. Negative thallium-201 SPECT/CT brain scan. Patient 1 (A), Patient 3 (B), Patient 4 (C), Patient 7 (D), Patient 8 (E), and Patient 9 (F) all have negative thallium-201 tracer uptake on planar and SPECT/CT imaging. SPECT/CT, single-photon emission computed tomography/computed tomography.