



Original Article

Corresponding Author

Tao Fan

https://orcid.org/0000-0001-8885-1480

Sanbo Brain Hospital, Capital Medical University, Beijing, People's Republic of China, NO.50 XiangshanYikesong Road, Haidian District, Beijing 100053, China
Email: fant@ccmu.edu.cn

Received: June 6, 2023

Revised: September 14, 2023

Accepted: September 16, 2023



This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<https://creativecommons.org/licenses/by-nc/4.0/>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

Copyright © 2023 by the Korean Spinal Neurosurgery Society

INTRODUCTION

Primary spinal cord astrocytoma is a rare disease with dismal functional and survival prognosis. In particular, patients with high-grade astrocytoma presented adverse overall survival (OS) time.^{1,2} Unlike its intracranial counterpart, high-grade intramedullary astrocytoma (grade, 3–4) lacks distinct boundaries with normal spinal cord tissue and does not have a unified treatment guideline.^{3–6} Furthermore, due to its rarity, prognostic factors associated with high-grade intramedullary astrocytoma and the role of H3K27M mutation in high-grade high-grade intramedullary astrocytoma have not been established.^{7–11} In

Does H3K27M Mutation Impact Survival Outcome of High-Grade Spinal Cord Astrocytoma?

Fan Zhang¹, Lei Cheng², Ze Ding¹, Shengxi Wang¹, Xingang Zhao¹, Zijun Zhao¹, Cong Liang¹, Kun Wu¹, Dongao Zhang¹, Yinqian Wang¹, Tao Fan¹

¹Sanbo Brain Hospital, Capital Medical University, Beijing, China

²Department of Neurosurgery, Xuanwu Hospital, China International Neuroscience Institute, Capital Medical University, Beijing, China

Objective: To evaluate the impact of H3K27M mutation in the prognosis of histological high-grade intramedullary astrocytoma.

Methods: A total of 78 patients who were diagnosed with high-grade spinal cord astrocytoma were included. Clinical data consisting demographic, radiological, molecular features and treatment data were recorded. Univariate and multivariate Cox analysis were performed to investigate variables associated with survival outcome of histological high-grade spinal cord astrocytoma.

Results: Median survival time was 21 months. Overall survival (OS) at 1 and 3 years was 65.7% and 40.7%, respectively. Sex, location, and tumor span did not present significant association with OS. Patients with H3K27M mutation showed significant shorter duration of symptom than patients with H3K27 wild-type. As respect to adjuvant treatment, adjuvant radiotherapy and chemotherapy were associated with favorable OS (both $p = 0.01$). Younger patients (age ≤ 18 years) had shorter OS ($p = 0.008$) than adult patients (age > 18 years). Of note, H3K27M mutation did not show significant impact on the survival outcome, regardless of histology grade 3 or grade 4 ($p = 0.3$).

Conclusion: Histological high-grade spinal cord astrocytoma has dismal prognosis. Our study demonstrated that H3K27M mutation did not show significant impact on survival outcome of histological high-grade spinal cord astrocytoma.

Keywords: Spinal cord, Astrocytoma, H3K27M, Oncology

this study, we aimed to evaluate the prognostic factors associated with high-grade intramedullary astrocytoma and, especially, the impact of H3K27M mutation on survival outcome of histological high-grade intramedullary astrocytoma in a large sample-size cohort.

MATERIALS AND METHODS

1. Patient Population

After reviewing medical records systems, a total of 66 patients with primary spinal cord astrocytoma between 2016 and 2021 at Beijing Sanbo Brain Hospital were retrieved. Among them,

13 cases and 5 cases were histologically classified as grade 1 and grade 2 spinal cord astrocytoma, respectively and were excluded. The remaining 48 patients who underwent surgery and were histologically diagnosed with high-grade spinal cord astrocytoma and included in our analysis. Clinical data, including age, sex, histological type, tumor location, preoperative modified McCormick Scale (MMS)³ (Supplementary Fig. 1), postoperative modified MMS, last follow-up modified MMS, extent of resection, and radiotherapy, are recorded. OS was calculated as the time from date of diagnosis to date of death or last follow-up time, while duration of symptoms was defined as the time from date of related-symptom onset to date of diagnosis.

Considering sample-size limitations, data from the literatures were also retrieved and information, including age, sex, the presence of H3K27M mutation, ATRX loss, high p53 expression, and Ki67 expression, were extracted (Table 1).¹²⁻¹⁴ Age groups were divided into pediatric (≤ 18 years) and adult group (> 18 years).^{13,15,16}

Our study is a investigator initiated trial, and the protocol of our study was approved by the Institutional Review Board of Capital Medical University (SBNK-YJ-2020-006-03). The informed written consent of the patient has been obtained.

2. Histological Morphology and Molecular Pathology

Firstly, histological morphology and grading were determined by 2 neuropathologists according to the 2021 World Health Organization (WHO) Classification.¹⁷⁻¹⁹ Tumors with anaplasia and mitotic activity would be defined as histological grade 3, while diffuse astrocytoma with microvascular proliferation and/or necrosis was designated as glioblastoma, histological grade 4. Secondly, status of H3K27M was determined using immunohistochemistry (IHC) method (positive H3K27M protein and loss of histone tri-methylation). Tumors with H3K27M mutation would be designated as diffuse midline glioma, WHO grade 4. Besides, ATRX loss, IDH1-R132H mutation, and p53 overexpression Ki-67 were also detected using IHC method. MGMT promotor methylation were detected by the Strepto-

myces antibiotic protein- peroxidase ligation method.²⁰

3. Treatment

All patients received surgery under intraoperative neurophysiological monitoring. The extent of resection was divided into gross total resection (GTR) ($\geq 90\%$), subtotal resection (SR) ($\geq 50\%$ and $< 90\%$), partial resection (PR) ($< 50\%$) and biopsy. Extent of resection was determined through operative magnetic resonance imaging (MRI) image (contrast-enhanced T1 weighted image) or determined by reviewing operation note and operative video. Illustrative cases were presented in Fig. 1. Postoperative adjuvant therapy was prescribed according to STUPP protocol.²¹ In detail, patients were given radiotherapy (1.8 Gy per daily fraction (Monday to Friday) over 6 weeks, total 45–50 Gy) plus concomitant temozolomide (75 mg/m², 7 days per week) and adjuvant temozolomide (6 cycles, 150 mg/m² for 5 days, 28 days cycle).

4. Statistical Analysis

The statistical analysis was performed using R ver. 4.2.1 (R Foundation for Statistical Computing, Vienna, Austria). Continuous variables were presented as mean $\pm$ standard deviations or median and interquartile ranges, while categorical variables were expressed as frequencies and proportions. Pearson chi-square test or Fisher exact test was used to compare categorical variables. Kaplan-Meier curves was constructed for survival by stratified variables and compared using log-rank test. Univariable and multivariable Cox proportional hazards regression analysis was used to determine risk factors associated with survival outcome. A p-value of < 0.05 was considered significant.

Table 1. Summary of recent literature on primary spinal cord astrocytoma

Study	Source	Year	Study design
Alvi et al. ¹⁴	<i>Mordon Pathology</i>	2019.06	Single center
Lebrun et al. ¹³	<i>Acta Neuropathologica Communications</i>	2020.08	Meta-analysis
Sloan et al. ¹²	<i>Acta Neuropathologica</i>	2019.01	Single institution




Fig. 1. Magnetic resonance imaging for illustrative cases.

Table 2. Clinical characteristics

Characteristic	Total (n = 78)	H3K27M WT (n = 34)	H3K27M mutant (n = 44)	p-value
Sex				0.25
Male	47 (60.3)	18 (52.9)	29 (65.9)	
Female	31 (39.7)	16 (47.1)	15 (34.1)	
Age (yr)				0.003
≤ 18	20 (25.6)	3 (8.8)	17 (38.6)	
> 18	58 (74.4)	31 (91.2)	27 (61.4)	
Histological grading				0.55
3	42 (53.8)	17 (50.0)	25 (56.8)	
4	36 (46.2)	17 (50.0)	19 (43.2)	
Duration of symptom (mo)				0.027
≤ 6	35 (44.9)	10 (29.4)	25 (56.8)	
> 6	13 (16.7)	9 (26.5)	4 (9.1)	
Unknown	30 (38.5)	15 (44.1)	15 (34.1)	
Site				0.70
Cervical	26 (33.3)	10 (29.4)	16 (36.4)	
Cervicothoracic	9 (11.5)	3 (8.8)	6 (13.6)	
Thoracic	30 (38.5)	16 (47.1)	14 (31.2)	
Thoracolumbar	7 (9.0)	4 (11.7)	3 (6.8)	
Lumbar	5 (6.4)	5 (14.7)	4 (9.1)	
Holocord	1 (1.3)	0 (0)	1 (2.3)	
Tumoral enhancement				0.21
None	7 (11.5)	1 (28.6)	6 (5.3)	
Partial	25 (34.6)	1 (14.3)	8 (42.1)	
Diffuse	18 (23.1)	4 (11.8)	14 (31.2)	
Unknown	28 (35.9)	12 (35.3)	16 (36.4)	
p53 overexpression				0.054
Yes	44 (31.3)	15 (44.1)	29 (65.9)	
No	34 (68.8)	19 (55.9)	15 (34.1)	
ATRX loss				0.90
Yes	30 (38.5)	14 (41.2)	16 (36.4)	
No	47 (60.3)	20 (58.8)	27 (61.4)	
Unknown	1 (1.3)	0 (0)	1 (2.3)	
Radiotherapy				0.33
Yes	52 (66.7)	23 (67.6)	29 (65.9)	
No	17 (21.8)	9 (26.5)	8 (18.2)	
Unknown	9 (11.5)	2 (5.9)	7 (15.9)	
Chemotherapy				0.42
Yes	53 (67.9)	26 (76.5)	27 (61.4)	
No	15 (19.2)	5 (14.7)	10 (22.7)	
Unknown	10 (12.9)	3 (8.8)	7 (15.9)	
Surgery				0.18
Biopsy	25 (32.1)	8 (23.5)	17 (38.6)	
Partial	32 (41.0)	8 (51.7)	15 (47.9)	
Gross total resection	21 (26.9)	12 (35.3)	9 (20.5)	

Values are presented as number (%).

WT, wild-type.

RESULTS

1. Clinical Data

A total of 78 patients who were histologically diagnosed as high-grade (grades 3–4) primary spinal cord astrocytoma based on tissue morphology were included. Out of these, 44 patients had H3K27M mutation and consequently, integrated diagnosis of the 44 patients was defined as diffuse midline glioma, H3K27M, WHO grade 4. Mean age of patients with and without H3K27M mutation was 29.4 and 42.2 years, respectively, and H3K27 wild-type (WT) frequently occurred in adult patients ($p = 0.003$). Forty-seven patients were male. The mean duration of symptom was 6.7 months, patients with H3K27M mutation exhibited shorter symptom duration than patients with H3K27 WT astrocytoma ($p = 0.027$) (Table 2).

2. Tumor Characteristics

For the location, the tumors were most frequently located in the thoracic cord (38.5%), followed by the cervical (33.3%), cervicothoracic (11.5%), and thoracolumbar (9.0%). The average length of tumor span was 4 segments. On contrast-enhanced MRI, partial enhancement was observed in 25 patients (34.6%), diffuse enhancement in 18 (23.1%), and no enhancement in 7 (11.5%). Image data for the remaining 28 patients were not available.

In terms of molecular features, none of the patients had IDH1-R132H mutation, while p53 overexpression occurred in 44 patients and ATRX loss occurred in 47 cases.

3. Treatment Strategy

Twenty-one patients underwent GTR ($\geq 90\%$), while biopsy and PR ($< 50\%$) were performed in 25 and 32 patients, respectively. A total of 57 patients received postoperative adjuvant treatment, out of which 48 patients received both chemotherapy and radiotherapy, while 4 patients received radiation therapy only and 5 patients received chemotherapy only, 10 patients did not receive adjuvant treatment and the postoperative treatment modality of the remaining 11 patients were unknown.

4. Survival Analysis

Median survival time was 21 months. OS at 1 and 3 years was 65.7% and 40.7%, respectively (Fig. 2). The median OS time of patients with and without H3K27M mutation was 15.3 months and 52.1 months respectively. We performed Kaplan-Meier curves and univariate and multivariate Cox analysis of clinicopathological features in 78 cases (Table 3). Kaplan-Meier curves




Fig. 2. Overall survival curve of 78 patients.

showed that sex, location and tumoral enhancement did not present significant association with OS (Fig. 3A, D, E). Patients with H3K27M mutation showed significant shorter duration of symptom than patients with H3K27 WT (Fig. 3C). As respect to adjuvant treatment, adjuvant radiotherapy and chemotherapy was associated with favorable OS (both $p = 0.01$) (Fig. 4A, B). Younger patients (age ≤ 18 years) had shorter OS ($p = 0.008$) than adult patients (age > 18 years) (Fig. 3B). Of note, patients with H3K27M mutation did not show worse survival outcome than patients with H3K27 WT ($p = 0.3$), regardless of histology grade 3 or grade 4 ($p = 0.9$, $p = 0.2$) (Fig. 5A–C). In terms of ATRX loss, our analysis did not reveal ATRX loss had a significant impact on survival outcome ($p = 0.07$) (Fig. 5D). Conversely, patients harboring p53 overexpression demonstrated a favorable OS compared to those normal p53 expression ($p = 0.009$) (Fig. 5E). In the context of neurological function, data were collected from 48 patients who underwent surgical treatment at Sanbo Brain Hospital. Among these, 6 patients underwent biopsy, 23 underwent PR, and 19 underwent gross total resection. Notably, the majority of patients (19 cases, 24.3%) exhibited MMS IV at the time of presentation. In contrast, 12 cases (15.3%) displayed modified MMS III, and 11 cases (14.1%) were classified as MMS V. A smaller subset of patients (3 cases, 3.8%) presented with mild neurological deficits (MMS I, MMS II), respectively. The

Table 3. Univariate and multivariate Cox analysis of clinicopathological features in spinal cord astrocytomas

Variable	Univariate		p-value	Multivariate		p-value
	HR	95% CI		HR	95% CI	
Histological grade	1.42	0.8–2.51	0.233	0.76	0.98–4.69	0.055
Surgery		-			-	
GTR	Reference	-	0.5	Reference	-	-
PR	0.74	0.39–1.4	0.352	0.28	0.09–0.82	0.02*
Biopsy	0.66	0.3–1.48	0.315	0.16	0.04–0.6	0.007*
H3K27M	1.726	0.95–3.15	0.08	0.83	0.37–1.87	0.659
Radiotherapy		-			-	
No	Reference	-	0.02*	Reference	-	-
Yes	0.37	0.17–0.79	0.01*	0.33	0.09–1.13	0.078
Unknown	0.90	0.35–2.3	0.829	0.04	0.002–0.63	0.023*
Chemotherapy		-			-	
No	Reference	-	0.02*	Reference	-	-
Yes	0.42	0.2–0.91	0.028*	0.62	0.15–2.49	0.504
Unknown	1.11	0.44–2.84	0.821	11.1	0.78–157.82	0.075
Sex	0.84	0.46–1.52	0.6		-	
Age	0.46	0.26–0.83	0.011*	0.33	0.14–0.76	0.009*
Site		-			-	
Cervical	Reference	-	0.1	Reference	-	-
Cervicothoracic	1.67	0.38–7.22	0.495	3.58	0.63–20.48	0.151
Thoracic	2.63	0.55–12.54	0.224	2.62	0.42–16.27	0.299
Thoracolumbar	0.82	0.18–3.7	0.799	1.57	0.28–8.9	0.609
Lumbar	1.05	0.2–5.58	0.954	1.17	0.18–7.63	0.865
Holocord	7.49	0.65–86.94	0.107	5.82	0.29–114.9	0.246
Duration of symptom		-			-	
≤ 6	Reference		0.06	Reference		-
> 6	0.52	0.21–1.26	0.15	1.01	0.32–3.23	0.983
Unknown	0.47	0.23–0.95	0.036*	0.56	0.15–2.11	0.396
Tumoral enhancement						
None	Reference		0.3	-		-
Partial	0.62	0.23–1.68	0.348	-	-	-
Diffuse	0.82	0.31–2.19	0.689	-	-	-
Unknown	0.47	0.18–1.23	0.125	-		-
p53 overexpression	2.23	1.2–4.137	0.01*	2.13	0.89–5.06	0.086
ATRX	0.54	0.28–1.05	0.069	0.63	0.26–1.5	0.292

HR, hazard ratio; CI, confidence interval; GTR, gross total resection; PR, partial resection.

*p < 0.05, statistically significant differences.

median preoperative modified MMS score was 4 (interquartile range [IQR], 3–4), while the postoperative MMS score was 5 (IQR, 4–5) (Fig. 6). Overall, no significant difference was observed between pre and postoperative MMS scores among different surgery groups ($p = 0.638$). In biopsy group, 4 cases exhibited postoperative worsening in their neurological status, 1 case

showed improvement, and 1 case displayed no change. In PR group, 12 cases experienced postoperative neurological deterioration, while 11 cases showed no change. In gross total resection group, 10 cases experienced postoperative functional decline, while 9 cases displayed no change. Furthermore, multivariable Cox analysis showed age > 18 (hazard ratio [HR], 0.33; 95% con-




Fig. 3. Kaplan-Meier survival curves of clinical features. (A) Sex. (B) Age. (C) Duration of symptom. (D) Tumor location. (E) Tumoral enhancement.




Fig. 4. The effect of treatment on survival outcome for patients with high-grade. (A) Radiotherapy. (B) Chemotherapy. (C) Surgery.




Fig. 5. Kaplan-Meier survival curves of molecular prognosis biomarker. (A) H3K27M mutation in high-grade spinal cord glioma. (B) H3K27M mutation in histological 3. (C) H3K27M mutation in histological 4. (D) ATRX. (E) p53.

fidence interval [CI], 0.14–0.76; $p = 0.009$), PR (HR, 0.28; 95% CI, 0.09–0.82; $p = 0.02$), and biopsy (HR, 0.16; 95% CI, 0.04–0.6; $p = 0.007$) were associated with favorable survival of high-grade spinal cord astrocytoma.

DISCUSSION

Primary spinal cord glioma represents 2% to 4% of all primary gliomas of the central nervous system, among which astrocytoma accounts for 2.4%–3%.^{1,22–24} H3K27M mutation was confirmed to be prognostic and diagnostic molecular biomarker of diffuse midline glioma, H3K27M-altered.^{13,20,24,25} Based on

previous reports, the presence of H3K27M mutation in grade 1 spinal cord astrocytoma has been seldom documented. The prevalence of the K27M mutation in grades 2, 3, and 4 varies across different studies. Some investigations indicate no significant susceptibility in the occurrence rate of the H3K27M mutation among grades 2, 3, and 4 tumors. For instance, Wang et al.²⁶ reported mutation rates of 36.1%, 39.3%, and 24.6% in histological grades 2, 3, and 4, respectively. Chai et al.²⁷ noted a mutation rate of 40% in both grades 2 and 3 tumors, contrasting with 20% in grade 4. Cheng et al.²⁸ documented mutation rates of 32.1% in grades 2 and 4, and 35.7% in grade 3, other studies have revealed a frequent occurrence of the H3K27M




Fig. 6. The neurofunction status before and after operative.

mutation in high-grade spinal cord astrocytoma.²⁹ Consistently, in our cohort, 40% (2 of 5), 59.5% (25 of 42), and 52.7% (19 of 36) in grade 2, 3, and 4 primary spinal cord astrocytomas harbored H3K27M mutation, respectively. Diffuse astrocytoma with H3K27M mutation was reported to have dismal survival outcome (median OS time, 20.7 months).²⁷ Of note, regardless of molecular biomarker, histological grading was always an essential prognostic factors and, similar to diffuse astrocytoma with H3K27M mutation, histological high-grade astrocytoma also showed poor prognosis.^{9,22,30,31} The impact of H3K27M mutation on histological high-grade spinal cord astrocytoma is unclear, does H3K27M mutation worsen poor prognosis of histological high-grade spinal cord astrocytoma? The study by Yi et al.⁷ revealed that patients with H3K27M mutant histological grade 4 astrocytoma had longer OS than patients with H3K27 WT histological grade 4 astrocytoma. However, Chai et al.²⁷ and Cheng et al.²⁸ found that there was no significant difference in prognosis between H3-WT and H3K27M-mutant histological grades 3–4 gliomas. Consistently, in our study, H3K27M mutation did not show association with survival outcome of histological high-grade spinal cord astrocytoma, regardless of histological grade 3 and grade 4 tumors. This finding likely suggested that histological grading was still the most important prognostic factor of primary spinal cord astrocytoma. However, small sample size of our study might limit the power to find

tiny prognostic difference potentially existed between histological high-grade astrocytoma with and without H3K27M mutation. A large sample-sized study is warranted to investigate the role of H3K27M mutation in histological high-grade intramedullary astrocytoma.

In term of treatment, the study by Wang et al. demonstrated that extent of resection and adjuvant treatment were not prognostic factors of OS in patients with primary spinal cord glioma.^{8,25,32} In a large cohort study, Chalif et al.³⁰ found that GTR and adjuvant therapy independently improve survival for high-grade spinal cord glioma. Our study which just focused on histological high-grade spinal cord glioma showed that GTR did not provide additional survival benefit, while patients could benefit from adjuvant radiotherapy or chemotherapy. GTR without function compromise for primary spinal cord glioma is a great challenge due to no definite margin existing between normal spinal cord and tumor. Consequently, adjuvant treatment was routinely prescribed to patients with high-grade spinal cord glioma in our center. Limitations of this study include the relatively small sample size, owing to the rarity of this disease. As a result, a detailed multivariate Cox regression analysis was no performed. Limited by the unavailability of sufficient samples for comprehensive analysis, only the fundamental molecular characteristics of the tumors were examined. Additionally, selection bias in treatment modality, which is one nature of retrospective study, might confound the final conclusion, hence, large sample-size prospective study is warranted to investigate the role of adjuvant treatment in primary spinal cord glioma.

CONCLUSION

Histological high-grade spinal cord astrocytoma has dismal prognosis and H3K27M mutation did not show significant impact on survival outcome of histological high-grade spinal cord astrocytoma.

NOTES

Supplementary Material: Supplementary Fig. 1 can be found via <https://doi.org/10.14245/ns.2346650.325>.

Conflict of Interest: The authors have nothing to disclose.

Funding/Support: This study received no specific grant from any funding agency in the public, commercial, or not-for-profit sectors.

Acknowledgments: We are particularly grateful to Jian Qiang and Cheng Jiahui for technical support.

Author Contribution: Conceptualization: LC, KW, DZ, TF; Data curation: FZ, YW; Formal analysis: DZ, XZ; Funding acquisition: CL, TF; Methodology: FZ, LC, SW, ZZ, KW, DZ, TF; Project administration: FZ, DZ, TF; Visualization: FZ, CL; Writing - original draft: FZ, LC, ZD, SW, ZZ, KW, YW; Writing - review & editing: XZ, CL, DZ, TF.

ORCID

Fan Zhang: 0009-0007-3163-5526

Lei Cheng: 0000-0001-6328-7852

Ze Ding: 0009-0000-5172-0719

Shengxi Wang: 0009-0007-5187-5869

Xingang Zhao: 0009-0005-8802-959X

Zijun Zhao: 0000-0002-9417-0224

Cong Liang: 0009-0004-3906-8237

Kun Wu: 0000-0002-6777-9413

Dongao Zhang: 0000-0002-0808-6203

Yinqian Wang: 0000-0001-5004-9417

Tao Fan: 0000-0001-8885-1480

REFERENCES

1. Chamberlain MC, Tredway TL. Adult primary intradural spinal cord tumors: a review. *Curr Neurol Neurosci Rep* 2011; 11:320-8.
2. Solomon DA, Wood MD, Tihan T, et al. Diffuse midline gliomas with histone H3-K27M mutation: a series of 47 cases assessing the spectrum of morphologic variation and associated genetic alterations. *Brain Pathol* 2016;26:569-80.
3. Manzano G, Green BA, Vanni S, et al. Contemporary management of adult intramedullary spinal tumors-pathology and neurological outcomes related to surgical resection. *Spinal Cord* 2008;46:540-6.
4. Shen G, Ma H, Pan L, et al. FDG PET/CT and MRI in primary spinal cord glioblastoma. *Clin Nucl Med* 2020;45:e144-e5.
5. Jiang T, Nam DH, Ram Z, et al. Clinical practice guidelines for the management of adult diffuse gliomas. *Cancer Lett* 2021;499:60-72.
6. López G, Oberheim Bush NA, Berger MS, et al. Diffuse non-midline glioma with H3F3A K27M mutation: a prognostic and treatment dilemma. *Acta Neuropathol Commun* 2017; 5:38.
7. Yi S, Choi S, Shin DA, et al. Impact of H3.3 K27M mutation on prognosis and survival of grade IV spinal cord glioma on the basis of new 2016 World Health Organization classification of the central nervous system. *Neurosurgery* 2019;84: 1072-81.
8. Chai RC, Yan H, An SY, et al. Genomic profiling and prognostic factors of H3 K27M-mutant spinal cord diffuse glioma. *Brain Pathol* 2023;33:e13153.
9. Argersinger DP, Rivas SR, Shah AH, et al. New developments in the pathogenesis, therapeutic targeting, and treatment of H3K27M-mutant diffuse midline glioma. *Cancers (Basel)* 2021;13:5280.
10. Raco A, Polli FM, Palmieri M, et al. Spinal corpectomy for the management of thoracic malignant intraspinal tumors in paraplegic or irreversibly, severely paraparetic patients: a technical remark. *J Clin Neurosci* 2019;68:308-11.
11. Mo Y, Duan S, Zhang X, et al. Epigenome programming by H3.3K27M mutation creates a dependence of pediatric glioma on SMARCA4. *Cancer Discov* 2022;12:2906-29.
12. Sloan EA, Cooney T, Oberheim Bush NA, et al. Recurrent non-canonical histone H3 mutations in spinal cord diffuse gliomas. *Acta Neuropathol* 2019;138:877-81.
13. Lebrun L, Meléndez B, Blanchard O, et al. Clinical, radiological and molecular characterization of intramedullary astrocytomas. *Acta Neuropathol Commun* 2020;8:128.
14. Alvi MA, Ida CM, Paolini MA, et al. Spinal cord high-grade infiltrating gliomas in adults: clinico-pathological and molecular evaluation. *Mod Pathol* 2019;32:1236-43.
15. Ludwig K, Kornblum HI. Molecular markers in glioma. *J Neurooncol* 2017;134:505-12.
16. Castel D, Philippe C, Calmon R, et al. Histone H3F3A and HIST1H3B K27M mutations define two subgroups of diffuse intrinsic pontine gliomas with different prognosis and phenotypes. *Acta Neuropathol* 2015;130:815-27.
17. Komori T. The 2016 WHO classification of tumours of the central nervous system: the major points of revision. *Neurol Med Chir (Tokyo)* 2017;57:301-11.
18. Villa C, Miquel C, Mosses D, et al. The 2016 World Health Organization classification of tumours of the central nervous system. *Presse Med* 2018;47(11-12 Pt 2):e187-200.
19. Horbinski C, Berger T, Packer RJ, et al. Clinical implications of the 2021 edition of the WHO classification of central nervous system tumours. *Nat Rev Neurol* 2022;18:515-29.
20. Sun P, Fan DJ, Fan T, et al. A prospective clinical study on MGMT protein expression and the effect of gene promoter methylation on sensitivity to chemotherapeutics in spinal glioma. *J Inflamm Res* 2021;14:4777-84.
21. Stupp R, Mason WP, van den Bent MJ, et al. Radiotherapy plus concomitant and adjuvant temozolomide for glioblastoma.

- toma. *N Engl J Med* 2005;352:987-96.
22. Guidetti B, Mercuri S, Vagnozzi R. Long-term results of the surgical treatment of 129 intramedullary spinal gliomas. *J Neurosurg* 1981;54:323-30.
 23. Ostrom QT, Cioffi G, Gittleman H, et al. CBTRUS statistical report: primary brain and other central nervous system tumors diagnosed in the United States in 2012-2016. *Neuro Oncol* 2019;21:v1-100.
 24. Santi M, Mena H, Wong K, et al. Spinal cord malignant astrocytomas. Clinicopathologic features in 36 cases. *Cancer* 2003;98:554-61.
 25. Wang YZ, Zhang YW, Liu WH, et al. Spinal cord diffuse midline gliomas With H3 K27m-mutant: clinicopathological features and prognosis. *Neurosurgery* 2021;89:300-7.
 26. Wang L, Li Z, Zhang M, et al. H3 K27M-mutant diffuse midline gliomas in different anatomical locations. *Hum Pathol* 2018;78:89-96.
 27. Chai RC, Zhang YW, Liu YQ, et al. The molecular characteristics of spinal cord gliomas with or without H3 K27M mutation. *Acta Neuropathol Commun* 2020;8:40.
 28. Cheng L, Wang L, Yao Q, et al. Clinicoradiological characteristics of primary spinal cord H3 K27M-mutant diffuse midline glioma. *J Neurosurg Spine*. 2021 Sep 24;1-12. doi: 10.3171/2021.4.SPINE2140. [Epub].
 29. Gessi M, Gielen GH, Dreschmann V, et al. High frequency of H3F3A (K27M) mutations characterizes pediatric and adult high-grade gliomas of the spinal cord. *Acta Neuropathol* 2015;130:435-7.
 30. Chalif EJ, Foster C, Sack K, et al. Impact of extent of resection and adjuvant therapy in diffuse gliomas of the spine. *Spine J* 2023;23:1015-27.
 31. Brat DJ, Aldape K, Colman H, et al. cIMPACT-NOW update 3: recommended diagnostic criteria for "Diffuse astrocytic glioma, IDH-wildtype, with molecular features of glioblastoma, WHO grade IV". *Acta Neuropathol* 2018;136:805-10.
 32. Yao J, Wang L, Ge H, et al. Diffuse midline glioma with H3 K27M mutation of the spinal cord: a series of 33 cases. *Neuropathology* 2021;41:183-90.

Modified McCormick scale

Grade	Modified McCormick scale
I	Intact neurologically, normal ambulation, minimal dyesthesia
II	Mild motor or sensory deficit, functional independence
III	Moderate deficit, limitation of function, independent w/external aid
IV	Severe motor or sensory deficit, limited function, dependent
V	Paraplegia or quadriplegia, even w/flickering movement

Supplementary Fig. 1. Modified McCormick scale.